arXiv:0806.1018v2 [gr-qc] 14 Aug 2008

Hamiltonian thermodynamics of three-dimensional dilatonic black holes

Gonçalo A. S. Dias*
Centro Multidisciplinar de Astrofísica - CENTRA
Departamento de Física,
Instituto Superior Técnico - IST,
Universidade Técnica de Lisboa - UTL,
Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal

José P. S. Lemos†
Centro Multidisciplinar de Astrofísica - CENTRA
Departamento de Física,
Instituto Superior Técnico - IST,
Universidade Técnica de Lisboa - UTL,
Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal

The action for a class of three-dimensional dilaton-gravity theories with a negative cosmological constant can be recast in a Brans-Dicke type action, with its free ω parameter. These theories have static spherically symmetric black holes. Those with well formulated asymptotics are studied through a Hamiltonian formalism, and their thermodynamical properties are found out. The theories studied are general relativity ($\omega \to \infty$), a dimensionally reduced cylindrical four-dimensional general relativity theory ($\omega = 0$), and a theory representing a class of theories ($\omega = -3$). The Hamiltonian formalism is setup in three dimensions through foliations on the right region of the Carter-Penrose diagram, with the bifurcation 1-sphere as the left boundary, and anti-de Sitter infinity as the right boundary. The metric functions on the foliated hypersurfaces are the canonical coordinates. The Hamiltonian action is written, the Hamiltonian being a sum of constraints. One finds a new action which yields an unconstrained theory with one pair of canonical coordinates $\{M, P_M\}$, M being the mass parameter and P_M its conjugate momenta The resulting Hamiltonian is a sum of boundary terms only. A quantization of the theory is performed. The Schrödinger evolution operator is constructed, the trace is taken, and the partition function of the canonical ensemble is obtained. The black hole entropies differ, in general, from the usual quarter of the horizon area due to the dilaton.

PACS numbers: 04.60.Ds, 04.20.Fy, 04.60.Gw, 04.60.Kz, 04.70.Dy

I. INTRODUCTION

A. Black hole thermodynamics

Hawking radiation [1] is a phenomenon that emerges when one combines classical general relativity with quantum fields in a black hole background. It is thus a semiclassical phenomenon. The radiation is thermal and emitted at a given temperature T which depends on the black hole parameters. When this is linked to other black hole properties one finds that black holes indeed have entropy S and allow a thermodynamical description [2].

B. Path integral approach to black hole thermodynamics

One can also get the thermodynamic properties of a black hole through a path integral approach to quantum gravity. In this approach one uses Feynman's idea that the amplitude $< g_2, \Sigma_2, t_2 | g_1, \Sigma_1, t_1 >$ to go to a bra vector state $< g_2, \Sigma_2, t_2 |$, with metric g_2 on a spatial hypersurface Σ_2 at some generic prescribed time t_2, from one ket vector state, $| g_1, \Sigma_1, t_1 >$, with metric g_1 on a spatial hypersurface Σ_1 at some generic prescribed time t_1, can be found from the sum of all possible classical vacuum geometries, i.e., metric field configurations g, which take the prescribed values g_1 and g_2 on the surfaces Σ_1 and Σ_2, respectively, i.e., $< g_2, \Sigma_2, t_2 | g_1, \Sigma_1, t_1 >= \int D[g] \exp(iS[g])$, where $D[g]$ is a measure on the space of all the gravitational field configurations, and S is the Lorentzian action of the theory. Now,

---

* Email: gadias@fisica.ist.utl.pt
† Email: lemos@fisica.ist.utl.pt

it is known that if one Euclideanizes the prescribed time t such that $t = -i\beta$ and Euclideanizes the action $I = -iS$, one obtains a timeless amplitude, which is equivalent to the partition function $Z = \int D[g]\, \exp\left(-S[g]\right)$. The partition function is then computed in the saddle point approximation, which finally connects to thermodynamics through the relation between Z and the Helmholtz free energy F, i.e., $F = -\mathbf{T} \ln Z$, where $\mathbf{T} = 1/\beta$ is the temperature, and we put Boltzmann's constant $k_{\rm B} = 1$. Since one has computed the partition function Z in this formalism one is working, a priori, in a well defined canonical ensemble, where the temperature $\mathbf{T}$ is given. However, solitary black holes are unstable against thermal fluctuations, indeed the heat capacity is in general negative. This means that the canonical ensemble is not defined at all and one should work instead in a microcanonical ensemble [3]. In the microcanonical ensemble the total energy E of the system, rather than the temperature $\mathbf{T}$, is given, and the ensemble is usually well defined. On the other hand, calculations in the canonical ensemble are much easier than in the microcanonical. Thus, in order to stick to the canonical ensemble in a coherent fashion, one has to go beyond a solitary black hole, by enclosing it in an appropriate finite box. In this way, the path integral formalism and its inherent connection to the canonical ensemble, makes sense, if one carefully chooses the boundary conditions by fixing the local temperature at the surface of the box itself, as was done for the Schwarzschild case in [4], and generalized to an arbitrary static field in [5]. This can also be naturally extended to the grand canonical ensemble when one includes charge [6]. If further, one puts a cosmological constant into the system, i.e., one deals with an asymptotic anti-de Sitter spacetime, one can dispense with the box, since the cosmological constant itself yields a natural box [7]. This approach, called the Brown-York formalism, has been further used a number of times in various different dimensions and for several different theories of gravity containing black holes. One should note that since the entropy and the other laws of black hole thermodynamics can be derived without considering matter fields, one finds that the black hole entropy is really an intrinsic entropy pertaining to the black hole geometry itself.

C. Hamiltonian approach to black hole thermodynamics

There are many other approaches to calculate the entropy and the thermodynamics of a black hole. The route we want to follow, motivated by the success of the path integral approach, is to build a Lorentzian Hamiltonian classical theory of the gravity in question, and then obtain a Lorentzian time evolution operator in the Schrödinger picture. Afterward one performs a Wick rotation from real to imaginary time, in order to find a well defined partition function. In more detail, the prescription implicit in this approach is: find the Hamiltonian of the system, calculate then the time evolution between a final state and an initial state, i.e., between the bra and ket vectors of those states, $< g_2, \Sigma_2, t_2|$ and $|g_1, \Sigma_1, t_1 >$, and then Euclideanize time. Here, the amplitude to propagate to a configuration $< g_2, \Sigma_2, t_2|$ from a configuration $|g_1, \Sigma_1, t_1 >$, is represented by $< g_2, \Sigma_2, t_2| \exp\left(-iH(t_2 - t_1)\right) |g_1, \Sigma_1, t_1 >$ in the Schrödinger picture. Euclideanizing time, $t_2 - t_1 = -i\beta$ and summing over a complete orthonormal basis of configurations g_n one obtains the partition function $Z = \sum \exp\left(-\beta E_n\right)$, of the field g at a temperature $1/\beta$, where E_n is the eigenenergy of the eigenstate g_n. This route is based on the Hamiltonian methods of [8, 9, 10, 11]. It was developed by Louko and Whiting in [12] for the specific problem of finding black hole entropies and thermodynamic properties, and further applied in [13, 14, 15, 16] by Louko and collaborators, in [17] by Bose and collaborators, and in [18, 19] by Kunstatter and collaborators. This approach also points to an entropy where the degrees of freedom are in the gravitational field itself, since nowhere one mentions matter fields.

The Louko-Whiting method [12] relies heavily on the Hamiltonian approach of [11], which in turn is an important ramification of the Arnowitt, Deser, and Misner approach, the ADM approach [9] (see also [8]), when applied to the full vacuum Schwarzschild black hole spacetime. This spacetime is better described by a spherically symmetric white hole plus black hole plus two asymptotically flat regions. These regions are well pictured in a Kruskal, or perhaps better, in a Carter-Penrose diagram. Indeed, Kuchař [11] in studying within this formalism the Schwarzschild black hole found the true dynamical degree of freedom of the phase space of such a spacetime, by considering the spacelike foliations of the full manifold. This degree of freedom is represented by one pair of canonical variables. This pair is composed of the mass M of the solution and its conjugate momentum, which physically represents the difference between the Killing times at right and left spatial infinities. In [12], this method was adapted by considering a spacelike foliation to the right of the future event horizon of the solution, enabling one to find the corresponding reduced phase space. From this one can obtain a Hamiltonian H and thus the Lorentzian time evolution operator $\exp(-iHt)$ in the Schrödinger picture.

This method has been applied for various theories of gravity and in several different dimensions. In [12], the first paper of the series, a vacuum Schwarzschild black hole in four dimensions in general relativity was placed in a box, defined as a rigid timelike frontier. The values of the metric functions were fixed both at the horizon and at the box. Then, in [13], the same procedure was applied to a vacuum dilatonic black hole in two dimensions. Here there was also a rigid frontier, where the values of the fields, dilatonic and gravitational, were fixed. Afterward, in [14], the method was applied to the Reissner-Nördstrom anti-de Sitter black hole in four-dimensional Einstein-Maxwell theory with a negative cosmological constant. This time there was no rigid frontier, so the far away asymptotic properties were defined at anti-de Sitter infinity. Next, in [15], the same formalism is applied to the case of topological black hole solutions of the equations of motion for the four-dimensional Einstein-Maxwell-anti-de Sitter theory, leaving however part of the Hamiltonian method implicit. Finally, it was shown in [16] that the entropy could also be calculated via this Hamiltonian method for five-dimensional spherically symmetric solutions of a one parameter family included within Lovelock gravity theory, where the action is comprised of the Ricci scalar term and the four-dimensional Euler density, i.e. the Gauss-Bonnet term, multiplied by an undetermined coefficient. There were some other developments. In [17] the Brown-York approach and the Louko-Whiting approach are compared. The main differences lie in the choice of boundary conditions, resulting in the fact that the Hamiltonian of Brown-York is the internal energy, whereas the Louko-Whiting Hamiltonian is the Helmholtz free energy. In [18, 19] the Louko-Whiting method was applied with some modifications to generic two-dimensional dilaton-gravity theories. Here we want to use the Louko-Whiting method to study the entropy and thermodynamic properties of three-dimensional black hole solutions in gravity theories with a dilaton, a negative cosmological constant and a free parameter ω, which can be generically comprised in a three-dimensional Brans-Dicke theory with a cosmological constant.

D. Three-dimensional dilaton-gravity black holes and their Hamiltonian approach to thermodynamics

There is great interest in studying three-dimensional theories of gravity. One of the main reasons is richness of structure with one dimension less [20, 21, 22, 23, 24, 25, 26, 27, 28]. In fact, it is simpler to deal with certain concepts such as temperature, entropy, and flux of radiation when one works in three dimensions, and it also seems easier to try a three-dimensional quantum description of the black hole system. Theories in three dimensions, such as general relativity, have a rich structure. Indeed, three-dimensional general relativity with a negative cosmological constant has a black hole solution, the Bañados-Teitelboim-Zanelli (BTZ) black hole [20, 21]. The BTZ black hole is also a solution of string theory with a dilaton and other fields [22], and there is also a three dimensional black string solution which can be recovered from an exact conformal field theory [23]. The existence of classical solutions depends on there being a negative cosmological constant. A negative constant is part of the three-dimensional black hole spacetimes structure, where the term in the action assumes a form given by $-2\lambda^2$. The cosmological constant term is therefore always negative, and thus, the black hole solutions of three-dimensional theories are usually asymptotically anti-de Sitter. Beyond three-dimensional general relativity and effective three-dimensional string gravity theories there are other interesting three-dimensional theories with a cosmological constant. One of these can be recovered from dimensional reduction of four-dimensional cylindrical general relativity, yielding a three-dimensional gravity with a dilaton with a particular coupling for the kinetic dilaton term [24]. It is then natural to set up a general three-dimensional dilaton-gravity theory, by including an ω paraneter, which yields different couplings for the kinetic term of the dilaton. This theory is then Brans-Dicke theory in three dimensions [25]. We will study in this connection the three-dimensional black hole solutions of [25], which are parametrized by the Brans-Dicke parameter ω, where, in principle, ω can take any value, i.e., $\infty > \omega > -\infty$. Since we want to perform a canonical Hamiltonian analysis, using an ADM formalism supplied with proper boundary conditions, it is necessary to pick up from the maze of solutions found in [25], only those that fulfill the boundary conditions we want to impose. First, we are interested only in solutions with horizons. Second, we want only solutions that are asymptotically anti-de Sitter at infinity. The cases of interest to be studied are then black holes for which $\omega \to \pm\infty$, $\infty > \omega > -1$, and $-\frac{3}{2} > \omega > -\infty$. As in [25] we choose three typical amenable cases where an analytical study can be done. These are $\omega \to \infty$ (or equivalently $\omega \to -\infty$), $\omega = 0$, and $\omega = -3$. The theory for which $\omega \to \infty$ is general relativity, and the solution is the BTZ black hole [20]. The theory for which $\omega = 0$ is equivalent to cylindrical four-dimensional general relativity and the corresponding black hole was found in [24]. The theory for which $\omega = -3$ is just a case of 3D Brans-Dicke theory, with a black hole solution that can be analyzed in this context [25]. If a quantum theory only makes sense if its classical form can be quantized by Hamiltonian methods, one should pick up only solutions which can be put consistently in a Hamiltonian form.

Thus, using the prescription for the canonical variables, we foliate, following ADM, the three-dimensional spacetime with spacelike hypersurfaces of equal time, whose right end is at the anti-de Sitter infinity, and whose left end is at the bifurcation circle of a non degenerate Killing horizon. These canonical variables are then transformed to another set of appropriate canonical variables, directly related to the physical parameters of the black hole classical solutions. In turn, these latter variables are reduced to the true degrees of freedom, which can be represented by one pair of new canonical variables, the mass M and its conjugate momentum P_M. All this construction is limited to the right static region of the Carter-Penrose diagram of spacetime. This reduced theory is then quantized canonically. From the quantum theory obtained, one builds a time evolution operator in the Schrödinger picture. Upon continuation of this operator to imaginary time, inserting it into two generic base state vectors and taking the trace one obtains the canonical partition function, appropriate to a canonical ensemble. Note one needs to pay attention to the boundary conditions of the ensemble when trying to build an appropriate one for the geometries of a quantum theory of gravity.

These have to respect the stability properties of the semiclassical approximations of this quantum theory. Here these suitable boundary conditions are ensured by the fact that the black hole solutions are asymptotically anti-de Sitter. Now, the partition function obtained, given the right conditions, is dominated by classical Euclidean solutions. Following [14] we choose a renormalized Hawking temperature, which is taken as a fixed quantity in the canonical ensemble, due to the fact that the Hawking temperature of the black hole goes to zero at infinity. The entropy of the system can then be found using this whole formalism. We note that the entropy of black holes in three-dimensional dilaton-gravity theories of the kind we study, have been worked out in [26] through a completely different approach, namely, using known properties of two-dimensional conformal field theories. As well, the entropy and thermodynamics of the BTZ black hole have been exhaustively studied using a number of different methods, see [27] for a first study and [28] for a review.

E. Structure of the paper

The structure of the paper is as follows. In Sec. II A we present the classical solutions of the three-dimensional dilatonic black holes, whose quantization through Hamiltonian methods we will perform. There is a free parameter ω for which we choose three different values, $\omega=\infty$, 0, -3, corresponding to the BTZ black hole, the dimensionally reduce four-dimensional cylindrical black hole, and a three-dimensional dilatonic black hole, respectively. In Sec. II B we introduce the spacetime foliation through which we will define the canonical coordinates and which will allow us to write the action as a sum of constraints multiplied by their respective Lagrange multipliers. Then follow three sections, Secs. III, IV, and V, where we develop the thermodynamic Hamiltonian formalism for the $\omega=\infty$, 0, -3 black holes, respectively. In each section, we first give the metric and the dilaton fields (subsection **A**). Then we replace in the action of the theory the foliation ansatz, with the canonical coordinates, and determine the conjugate momenta of the coordinates. From there we determine the form of the Hamiltonian action through a Legendre transformation. It is then necessary to guarantee that there is a well defined variational principle, so as to allow the derivation of the equations of motion of the coordinates and respective momenta. This entails adding surface terms to the original action, which in turn forces the definition of asymptotic properties for the coordinates, momenta, and Lagrange multipliers. It is also necessary to fix some variables at the boundaries in order to have a well defined variational principle. After this is accomplished, we write the usual metric functions of a three-dimensional spherically symmetric, static black hole spacetime as functions of the canonical variables (subsection **B**). We then choose a new set of canonical variables and perform a canonical transformation. This new set has a physical meaning, i.e., one is the mass parameter of the black hole, and the other, the momentum conjugate to the mass, is the spatial derivative of the Killing time. We rewrite the action with the new variables and obtain a new pair of constraints with a redefined pair of Lagrange multipliers (subsection **C**). The new action is then reduced to its true degrees of freedom, with one pair of canonical variables, the mass and its conjugate momentum. The reduced Hamiltonian is nothing more than the initial surface terms written as functions of the new variables (subsection **D**). We quantize the theory canonically, defining a Hilbert space and an inner product therein, with the Hamiltonian now being an operator in this Hilbert space. We write the time evolution operator in the Schrödinger picture (subsection **E**). In order to obtain a thermodynamic description of the system we build a partition function in the canonical ensemble with fixed temperature by continuing the time evolution operator to imaginary time, and taking the trace in the mass eigenstates. This requires that we normalize the trace. Finally, through saddle point methods we obtain the thermodynamic functions (subsection **F**). In Sec. VI we discuss and conclude. Throughout the paper we make $\hbar=1$, $k_{\rm B}=1$, $c=1$, and $G=\frac{1}{8}$.

II. THE 3D BLACK HOLE SOLUTIONS WHICH ALLOW A PROPER THERMODYNAMIC HAMILTONIAN DESCRIPTION

A. The 3D black hole solutions

Three-dimensional black hole theories have been studied in, e.g., [20, 21, 22, 23, 24, 25, 26, 27, 28]. A general action that incorporates most of these black holes is a Brans-Dicke action, with gravitational and dilaton fields and a cosmological constant. It is given by [25]

$$S=\frac{1}{2\pi}\int d^3x\,\sqrt{-g}\,e^{-2\phi}\left(R-4\omega(\partial\phi)^2+4\lambda^2\right)+\bar{B}\,, \tag{1}$$

where g is the determinant of the three-dimensional metric $g_{\mu\nu}$, R is the curvature scalar, ϕ is a scalar dilaton field, λ is the cosmological constant, ω is the Brans-Dicke parameter, and $\bar{B}$ is a generic surface term.

The general solution for a static spherically symmetric metric, i.e., circularly symmetric (since we deal with tree dimensions), is [25]

$$
\begin{aligned}
ds^2 &= -\left[(aR)^2 - \frac{b}{(aR)^{\frac{1}{\omega+1}}}\right] dT^2 + \frac{dR^2}{(aR)^2 - \frac{b}{(aR)^{\frac{1}{\omega+1}}}} + R^2 d\varphi^2, & \omega \neq -2\,, -\frac{3}{2}\,, -1\,, && (2)\\
ds^2 &= -(R^2 - R)\, dT^2 + (R^2 - R)^{-1} dR^2 + R^2 d\varphi^2, & \omega = -2\,, && (3)\\
ds^2 &= 4\lambda^2 R^2 \ln(bR) dT^2 - \frac{dR^2}{4\lambda^2 R^2 \ln(bR)} R^2 d\varphi^2, & \omega = -\frac{3}{2}\,, && (4)\\
ds^2 &= -dT^2 + dR^2 + d\varphi^2, & \omega = -1, && (5)
\end{aligned}
$$

where T, R are Schwarzschild coordinates, a is a constant related to the cosmological constant (see below), and b is a constant of integration (see below), and the general solution for ϕ is given by

$$
\begin{aligned}
\phi &= -\frac{1}{2(\omega+1)} \ln(a\,R), & \omega \neq -1\,, && (6)\\
\phi &= \text{constant}\,, & \omega = -1\,. && (7)
\end{aligned}
$$

For the constant a one has

$$
\begin{aligned}
a &= \frac{2\,|(\omega+1)\lambda|}{\sqrt{|(\omega+2)(2\omega+3)|}}\,, & \omega \neq -2\,, -\frac{3}{2}\,, -1\,, && (8)\\
a &= 1\,, & \omega = -2, -\frac{3}{2}\,, && (9)\\
a &= 0\,, & \omega = -1\,. && (10)
\end{aligned}
$$

The constant b is related to the ADM mass of the solutions by

$$
\begin{aligned}
M &= \frac{\omega+2}{\omega+1}\, b\,, & \omega \neq -2\,, -\frac{3}{2}\,, -1\,, && (11)\\
M &= 0, & \omega = -2,\ -1, && (12)\\
M &= -4\lambda^2 \ln b, & \omega = -\frac{3}{2}\,. && (13)
\end{aligned}
$$

Since we want to perform a canonical Hamiltonian analysis, using an ADM formalism supplied with proper boundary conditions, it is necessary to pick up only those solutions that fulfill the conditions we want to impose. First, we are interested only in solutions with horizons, so we take b to be positive. Second, apart from a measure zero of solutions, all solutions have a non-zero $|\lambda|$. This does not mean straight away that the solutions are asymptotically anti-de Sitter. Some have one type or another of singularities at infinity, which do not allow an imposition of proper boundary conditions. So, from [25] with the corresponding Carter-Penrose diagrams, we discard the following solutions: $\omega = -1$ which is simply the Minkowski solution of a low-energy limit of string theory, $-1 > \omega > -\frac{3}{2}$ since it gives weird conical singularities at Carter-Penrose infinity, and $\omega = -\frac{3}{2}$ since all the Carter-Penrose boundary is singular. Thus, the cases of interest to be studied are black holes for which $\omega \to \infty$, $\infty > \omega > -1$, and $-\frac{3}{2} > \omega > -\infty$. For b positive these solutions have ADM mass M positive, so well defined horizons. As in [25] we choose three typical amenable cases where an analytical study can be done. These are $\omega \to \infty$ (or equivalently $\omega \to -\infty$), $\omega = 0$, and $\omega = -3$. The theory for which $\omega \to \infty$ is general relativity, and the solution is the BTZ black hole [20, 21]. The theory for which $\omega = 0$ is equivalent to cylindrical four-dimensional general relativity and the corresponding black hole was found in [24]. The theory for which $\omega = -3$ is just a case of 3D Brans-Dicke theory, with a particular form for the kinetic dilaton term, which has a black hole solution that can be analyzed in this context [25].

B. ADM form of the metric

The ansatz for the metric and dilaton fields with which we start our canonical analysis is given by

$$
ds^2 = -N(t,r)^2 dt^2 + \Lambda(t,r)^2 (dr + N^r(t,r)dt)^2 + R(t,r)^2 d\varphi^2\,, \tag{14}
$$

$$
\mathrm{e}^{-2\phi} = (a\,R(t,r))^{\frac{1}{\omega+1}}\ . \tag{15}
$$

This is the ADM ansatz [9] for the metric of spherically symmetric solutions of the three-dimensional Brans-Dicke theory. In this we follow the basic formalism developed by Kuchař [11]. The canonical coordinates R, Λ are functions of t and r, $R = R(t,r)$, $\Lambda = \Lambda(t,r)$. Now, $r = 0$ is generically on the horizon as analyzed in [11], but for our purposes $r = 0$ represents the horizon bifurcation point of the Carter-Penrose diagram [12] (see also [13]-[17]). In three spacetime dimensions the point represents a circle. The coordinate r tends to ∞ as the coordinates themselves tend to infinity, and t is another time coordinate. The remaining functions are the lapse $N = N(t,r)$ and shift functions $N^r = N^r(t,r)$ and will play the role of Lagrange multipliers of the Hamiltonian of the theory. The canonical coordinates $R = R(t,r)$, $\Lambda = \Lambda(t,r)$ and the lapse function $N = N(t,r)$ are taken to be positive. The angular coordinate is left untouched, due to spherical (i.e, circular) symmetry. The dilaton is a simple function of the radial canonical coordinate, and it can be traded directly by it through equation (15), as will be done below. The ansatz (14)-(15) is written in order to perform the foliation of spacetime into spacelike hypersurfaces, and thus separates the spatial part of the spacetime from the temporal part. Indeed, the canonical analysis requires the explicit separation of the time coordinate from the other space coordinates, and so in all expressions time is treated separately from the other coordinates. It breaks explicit, but not implicit, covariance of the three-dimensional Brans-Dicke theory. Such a split is necessary in order to perform the Hamiltonian analysis. The metric coefficients of the induced metric on the hypersurfaces become the canonical variables, and the momenta are determined in the usual way, by replacing the time derivatives of the canonical variables, the velocities. Then, using the Hamiltonian one builds a time evolution operator to construct an appropriate thermodynamic ensemble for the geometries of a quantum theory of gravity. Assuming that a quantum theory only makes sense if its classical form can be quantized by Hamiltonian methods, one should pick up only solutions which can be put consistently in a Hamiltonian form. Thus, in the following we perform a Hamiltonian analysis to extract the entropy and other thermodynamic properties in the three three-dimensional Brans-Dicke black holes mentioned above, those for which $\omega = \infty, 0, -3$.

III. HAMILTONIAN THERMODYNAMICS OF THE BTZ BLACK HOLE ($\omega = \infty$)

A. The metric

For $\omega \to \infty$, the three-dimensional Brans-Dicke theory reduces to three-dimensional general relativity [25]. Then the general metric and dilaton solutions, given in Eqs. (2) and (6), reduce to the following

$$ds^2 = -\left[(a\,R)^2 - M\right]\, dT^2 + \frac{dR^2}{(a\,R)^2 - M} + R^2\, d\varphi^2\,, \tag{16}$$

$$e^{-2\phi} = 1\,, \tag{17}$$

with $M = b$ and $a = \sqrt{2}|\lambda|$. This is the BTZ black hole solution. Next, in Fig. 1, we have the Carter-Penrose diagram of the BTZ black hole, where $R_{\rm h}$ denotes the black hole horizon radius, $R_{\rm h} = \sqrt{M/(2\lambda^2)}$, $R = 0$ is the radius of the causal singularity, and $R = \infty$ is the spatial infinity, and we have discriminated the static and dynamical regions by roman numerals, namely the static right I and left I', and the dynamical future II and past II' regions.



FIG. 1: The Carter-Penrose Diagram for the $\omega = \infty$ case.

B. Canonical formalism

The action (1) with $\omega \to \infty$, and $2\lambda^2 = l^{-2}$, where l is the AdS length, becomes, excluding surface terms,

$$\begin{aligned} S[\Lambda,\, R,\, \dot\Lambda,\, \dot R;\, N,\, N^r] \;=\; \int dt \int_0^\infty dr \Big\{ & -2N^{-1}\dot\Lambda\dot R + 2N^{-1}N^r R'\dot\Lambda + 2N^{-1}(N^r)'\Lambda\dot R + 2N^{-1}N^r\Lambda'\dot R \\ & -2N^{-1}(N^r)^2\Lambda' R' + 2N\Lambda^{-2}\Lambda' R' - 2N\Lambda^{-1}R'' + 4N\lambda^2\Lambda R \Big\}\,, \end{aligned} \qquad (18)$$

where $\dot{}$ means derivative with respect to time t and $'$ is the derivative with respect to r, and where all the explicit functional dependences are omitted. There are two ways to arrive at the simple form of Eq. (18). One is to replace the ADM ansatz for the metric and the field ϕ, given in (14)-(15), into the action (1). One obtains an expanded action, where one has polynomials on the canonical coordinates and the Lagrange multipliers, plus their derivatives, with respect both to time t and to space r. Afterward, one integrates in the angular coordinate φ and obtains a two-dimensional action. Of the terms composing it some are total derivatives which can be discarded. This may prove itself to be cumbersome. The other, simpler, way to arrive at the form of the action (18), is to dimensionally reduce, through the Killing angle coordinate φ, the action before replacing the ansatz (14)-(15). The new action is then a two-dimensional integral. In this way the total derivatives are seen straight away, allowing one to discard all the unnecessary surface terms. After the reduction and integration one can replace again the ansatz. Depending on the situation we use three different letters containing the same information but with slightly different numerical values. Thus, a, λ, and l are related by $a^2 = 2\lambda^2 = l^{-2}$, where l is the AdS length. From the action (18) one determines the canonical momenta, conjugate to Λ and R, respectively,

$$P_\Lambda \;=\; -2N^{-1}\left\{\dot R - R'N^r\right\}\,, \qquad (19)$$

$$P_R \;=\; -2N^{-1}\left\{\dot\Lambda - (\Lambda N^r)'\right\}\,. \qquad (20)$$

By performing a Legendre transformation, we obtain

$$\begin{aligned} \mathcal{H} \;=\; & N\left\{-\frac{1}{2}P_R P_\Lambda - 2\Lambda^{-2}\Lambda' R' + 2\Lambda^{-1}R'' - 4\lambda^2\Lambda R\right\} + N^r\left\{P_R R' - P_\Lambda'\Lambda\right\} \\ \equiv\; & NH + N^r H_r\,. \end{aligned} \qquad (21)$$

Here additional surface terms have been ignored, as for now we are interested in the bulk terms only. The action in Hamiltonian form is then

$$S[\Lambda,\, R,\, P_\Lambda,\, P_R;\, N,\, N^r] = \int dt \int_0^\infty dr \left\{P_\Lambda\dot\Lambda + P_R\dot R - NH - N^r H_r\right\}\,. \qquad (22)$$

The equations of motion are

$$\dot\Lambda \;=\; -\frac{1}{2}NP_R + (N^r\Lambda)'\,, \qquad (23)$$

$$\dot R \;=\; -\frac{1}{2}NP_\Lambda - N^r R'\,, \qquad (24)$$

$$\dot P_R \;=\; 4\lambda^2 N\Lambda - (2N'\Lambda^{-1})' + (N^r P_R)'\,, \qquad (25)$$

$$\dot P_\Lambda \;=\; 4\lambda^2 NR - 2N'\Lambda^{-2}R' + N^r(P_\Lambda)'\,. \qquad (26)$$

In order to have a well defined variational principle, we need to eliminate the surface terms of the original bulk action, which render the original action itself ill defined when one seeks a correct determination of the equations of motion through variational methods. These surface terms are eliminated through judicious choice of extra surface terms which should be added to the action. The action (22) has the following extra surface terms, after variation

$$\text{Surface terms} = \left(-N^r P_R\delta R + N^r\Lambda\delta P_\Lambda + 2N\Lambda^{-2}R'\delta\Lambda - 2N\Lambda^{-1}\delta R' + 2N'\Lambda^{-1}\delta R\right)\Big|_0^\infty\,. \qquad (27)$$

In order to evaluate this expression, we need to know the asymptotic conditions of each of the above functions individually, which are functions of (t, r).

Starting with the limit $r \to 0$ we assume

$$\Lambda(t, r) \;=\; \Lambda_0 + O(r^2)\,, \qquad (28)$$

$$\begin{aligned} R(t,r) &= R_0 + R_2 r^2 + O(r^4)\,, & (29)\\ P_\Lambda(t,r) &= O(r^3)\,, & (30)\\ P_R(t,r) &= O(r)\,, & (31)\\ N(t,r) &= N_1(t) r + O(r^3)\,, & (32)\\ N^r(t,r) &= O(r^3)\,. & (33)\end{aligned}$$

With these conditions, we have for the surface terms at $r=0$

$$\text{Surface terms}\big|_{r=0} = -2N_1\Lambda_0^{-1}\delta R_0\,. \tag{34}$$

Note that there are time dependences on the left hand side of the falloff conditions and that there are no such dependences on the right hand side, in the lower orders of the expansion in r. This apparent discrepancy stems from the fact that there is in fact no time dependence in the lower orders of the majority of the functions, but there may still exist such dependence for higher orders. Nevertheless, terms such as R_0 are functions, independent of (t,r), thus constant, but undetermined. Their variation makes sense, as we may still vary between different values for these constant functions.

For $r \to \infty$ we have

$$\begin{aligned} \Lambda(t,r) &= l r^{-1} + l^3 \eta(t) r^{-3} + O^\infty(r^{-5})\,, & (35)\\ R(t,r) &= r + l^2\rho(t) r^{-1} + O^\infty(r^{-3})\,, & (36)\\ P_\Lambda(t,r) &= O^\infty(r^{-2}) & (37)\\ P_R(t,r) &= O^\infty(r^{-4})\,, & (38)\\ N(t,r) &= R(t,r)'\Lambda(t,r)^{-1}(\tilde{N}_+(t) + O^\infty(r^{-5}))\,, & (39)\\ N^r(t,r) &= O^\infty(r^{-2})\,. & (40)\end{aligned}$$

Here, as usual, $l^{-2} = 2\lambda^2$. These conditions imply for the surface terms in the limit $r \to \infty$

$$\text{Surface terms}\big|_{r\to\infty} = 2\delta(M_+(t))\tilde{N}_+\,. \tag{41}$$

where $M_+(t) = 2(\eta(t) + 2\rho(t))$. So, the surface term added to (22) is

$$S_{\partial\Sigma}\left[\Lambda, R; N\right] = \int dt\left(2R_0 N_1\Lambda_0^{-1} - \tilde{N}_+ M_+\right)\,. \tag{42}$$

What is left after varying this last surface term and adding it to the varied initial action (see Eq. (22)) is

$$\int dt\left(2R_0\delta(N_1\Lambda_0^{-1}) - \delta\tilde{N}_+ M_+\right)\,. \tag{43}$$

We choose to fix $N_1\Lambda_0^{-1}$ on the horizon ($r=0$) and $\tilde{N}_+$ at infinity. These choices make the surface variation (43) disappear. The term $N_1\Lambda_0^{-1}$ is the integrand of

$$n^a(t_1)n_a(t_2) = -\cosh\left(\int_{t_1}^{t_2} dt\, N_1(t)\Lambda_0^{-1}(t)\right)\,, \tag{44}$$

which is the rate of the boost suffered by the future unit normal to the constant t hypersurfaces defined at the bifurcation circle, i.e., at $r \to 0$, due to the evolution of the constant t hypersurfaces. By fixing the integrand we are fixing the rate of the boost, which allows us to control the metric singularity when $r \to 0$ [12].

C. Reconstruction, canonical transformation, and action

In order to reconstruct the mass and the time from the canonical data, which amounts to making a canonical transformation, we have to rewrite the form of the solutions of Eqs. (16)-(17). We follow Kuchař [11] for this reconstruction. We concentrate our analysis on the right static region of the Carter-Penrose diagram. In the static region, we define F as

$$F(R(t,r)) = (aR(t,r))^2 - b\,, \tag{45}$$

and make the following substitutions

$$T = T(t,r)\,, \qquad\qquad R = R(t,r)\,, \tag{46}$$

into the solution (16), getting

$$ds^2 \;=\; -(F\dot{T}^2 - F^{-1}\dot{R}^2)\,dt^2 \;+\; 2(-FT'\dot{T} + F^{-1}R'\dot{R})\,dtdr \;+\; (-F(T')^2 + F^{-1}\dot{R}^2)\,dr^2 \;+\; R^2 d\varphi^2\,. \tag{47}$$

This introduces the ADM foliation directly into the solutions. Comparing it with the ADM metric (14), written in another form as

$$ds^2 = -(N^2 - \Lambda^2(N^r)^2)\,dt^2 \;+\; 2\Lambda^2 N^r\,dtdr \;+\; \Lambda^2 dr^2 \;+\; R^2\,d\varphi^2\,, \tag{48}$$

we can write a set of three equations

$$\begin{aligned} \Lambda^2 &= -F(T')^2 + F^{-1}(R')^2\,, & (49)\\ \Lambda^2 N^r &= -FT'\dot{T} + F^{-1}R'\dot{R}\,, & (50)\\ N^2 - \Lambda^2(N^r)^2 &= F\dot{T}^2 - F^{-1}\dot{R}^2\,. & (51) \end{aligned}$$

The first two equations, Eqs. (49) and Eq. (50), give

$$N^r = \frac{-FT'\dot{T} + F^{-1}R'\dot{R}}{-F(T')^2 + F^{-1}(R')^2}\,. \tag{52}$$

This one solution, together with Eq. (49), give

$$N = \frac{R'\dot{T} - T'\dot{R}}{\sqrt{-F(T')^2 + F^{-1}(R')^2}}\,. \tag{53}$$

One can show that $N(t,r)$ is positive (see [11]). Next, putting Eqs. (52)-(53), into the definition of the conjugate momentum of the canonical coordinate Λ, given in Eq. (19), one finds the spatial derivative of $T(t,r)$ as a function of the canonical coordinates, i.e.,

$$-T' = \frac{1}{2}F^{-1}\Lambda P_\Lambda\,. \tag{54}$$

Later we will see that $-T' = P_M$, as it will be conjugate to a new canonical coordinate M. Following this procedure to the end, we may then find the form of the new coordinate $M(t,r)$, as a function of t and r. First, we need to know the form of F as a function of the canonical pair Λ, R. For that, we replace back into Eq. (49) the definition, in Eq. (54), of T', giving

$$F = \left(\frac{R'}{\Lambda}\right)^2 - \left(\frac{P_\Lambda}{2}\right)^2\,. \tag{55}$$

Equating this form of F with Eq. (45), we obtain

$$M = 2\lambda^2 R^2 - F\,, \tag{56}$$

where F is given in Eq. (55) and $a^2 = 2\lambda^2$, see Eq. (8). We thus have found the form of the new canonical coordinate, M. It is now a straightforward calculation to determine the Poisson bracket of this variable with $P_M = -T'$ and see that they are conjugate, thus making Eq. (54) the conjugate momentum of M, i.e.,

$$P_M = \frac{1}{2}F^{-1}\Lambda P_\Lambda\,. \tag{57}$$

It is now necessary to find out the other new canonical variable which commutes with M and P_M and which guarantees, with its conjugate momentum, that the transformation from Λ, R, to M and the new variable is canonical. Immediately is it seen that R commutes with M and P_M. It is then a candidate. It remains to be seen whether P_R also commutes with M and P_M. As with R, it is straightforward to see that P_R does not commute with M and P_M, as these contain powers of R in their definitions, and $\{R(t,r),\, P_R(t,r^*)\} = \delta(r - r^*)$. So rename the canonical variable R as $R = \mathrm{R}$. We have then to find a new conjugate momentum to R which also commutes with M and P_M, making the transformation from $\{\Lambda,\, R;\, P_\Lambda,\, P_R\} \to \left\{M,\, \mathrm{R};\, P_M,\, P_{\mathrm{R}}\right\}$ a canonical one. The way to proceed is to look at the constraint H_r, which is called in this formalism the super-momentum. This is the constraint which generates spatial diffeomorphisms in all variables. Its form, in the initial canonical coordinates, is $H_r = -\Lambda\, P'_\Lambda + P_R\, R'$. In this formulation, Λ is a spatial density and R is a spatial scalar. As the new variables, M and R, are spatial scalars, the generator of spatial diffeomorphisms is written as $H_r = P_M M' + P_{\mathrm{R}}\mathrm{R}'$, regardless of the particular form of the canonical coordinate transformation. It is thus equating these two expressions of the super-momentum H_r, with M and P_M written as functions of Λ, R and their respective momenta, that gives us the equation for the new P_{R}. This results in

$$\begin{aligned} P_{\mathrm{R}} &= P_R - 2\lambda^2 F^{-1}\Lambda P_\Lambda R + F^{-1}\Lambda^{-1}P_\Lambda R'' - F^{-1}\Lambda^{-2}P_\Lambda\Lambda' R' \\ &\quad - F^{-1}\Lambda^{-1}P'_\Lambda R' \,. \end{aligned} \tag{58}$$

We have now all the canonical variables of the new set determined. For completeness and future use, we write the inverse transformation for Λ and P_Λ,

$$\Lambda = \left((\mathrm{R}')^2 F^{-1} - P_M^2 F\right)^{\frac{1}{2}} \,, \tag{59}$$

$$P_\Lambda = 2FP_M\left((\mathrm{R}')^2 F^{-1} - P_M^2 F\right)^{-\frac{1}{2}} \,. \tag{60}$$

In summary, the canonical transformations are the following,

$$\begin{aligned} R &= \mathrm{R}\,, \\ M &= 2\lambda^2 R^2 - F\,, \\ P_{\mathrm{R}} &= P_R - 2\lambda^2 F^{-1}\Lambda P_\Lambda R + F^{-1}\Lambda^{-1}P_\Lambda R'' - F^{-1}\Lambda^{-2}P_\Lambda\Lambda' R' \\ &\quad - F^{-1}\Lambda^{-1}P'_\Lambda R'\,, \\ P_M &= \frac{1}{2}F^{-1}\Lambda P_\Lambda \,. \end{aligned} \tag{61}$$

It remains to be seen that this set of transformations is in fact canonical. In order to prove that the set of equalities in expression (61) is canonical we start with the equality

$$\begin{aligned} P_\Lambda\delta\Lambda + P_R\delta R - P_M\delta M - P_{\mathrm{R}}\delta\mathrm{R} &= \left(\delta R \ln\left|\frac{2R' + \Lambda P_\Lambda}{2R' - \Lambda P_\Lambda}\right|\right)' + \\ &\quad + \delta\left(\Lambda P_\Lambda + R'\ln\left|\frac{2R' - \Lambda P_\Lambda}{2R' + \Lambda P_\Lambda}\right|\right) \,. \end{aligned} \tag{62}$$

We now integrate expression (62) in r, in the interval from $r = 0$ to $r = \infty$. The first term on the right hand side of Eq. (62) vanishes due to the falloff conditions (see Eqs. (28)-(33) and Eqs. (35)-(40)). We then obtain the following expression

$$\int_0^\infty dr\, \left(P_\Lambda\delta\Lambda + P_R\delta R\right) - \int_0^\infty dr\, \left(P_M\delta M + P_{\mathrm{R}}\delta\mathrm{R}\right) = \delta\omega\,[\Lambda,\, R,\, P_\Lambda] \,, \tag{63}$$

where $\delta\omega\,[\Lambda,\, R,\, P_\Lambda]$ is a well defined functional, which is also an exact form. This equality shows that the difference between the Liouville form of $\{R,\, \Lambda;\, P_R,\, P_\Lambda\}$ and the Liouville form of $\left\{\mathrm{R},\, M;\, P_{\mathrm{R}},\, P_M\right\}$ is an exact form, which implies that the transformation of variables given by the set of equations (61) is canonical.

Armed with the certainty of the canonicity of the new variables, we can write the asymptotic form of the canonical variables and of the metric function $F(t,r)$. These are, for $r \to 0$

$$F(t,r) = 4R_2^2\Lambda_0^{-2}r^2 + O(r^4)\,, \tag{64}$$

$$\mathrm{R}(t,r) = R_0 + R_2\, r^2 + O(r^4)\,, \tag{65}$$

$$M(t,r) = 2\lambda^2 R_0^2 + \left(4\lambda^2 R_0 R_2 - 4R_2^2\Lambda_0^{-2}\right) r^2 + O(r^4)\,, \tag{66}$$

$$P_{\mathrm{R}}(t,r) = O(r)\,, \tag{67}$$

$$P_M(t,r) = O(r)\,. \tag{68}$$

For $r \to \infty$, we have

$$F(t,r) = 2\lambda^2\, r^2 - 2(\eta(t) + 2\rho(t)) + O^\infty(r^{-2})\,, \tag{69}$$

$$\begin{aligned} \mathrm{R}(t,r) &= r + (2\lambda^2)^{-1}\rho(t)\, r^{-1} + O^\infty(r^{-3})\,, && (70)\\ M(t,r) &= M_+(t) + O^\infty(r^{-2})\,, && (71)\\ P_{\mathrm{R}}(t,r) &= O^\infty(r^{-4})\,, && (72)\\ P_M(t,r) &= O^\infty(r^{-5})\,, && (73) \end{aligned}$$

where $M_+(t) = 2(\eta(t) + 2\rho(t))$, as seen before in the surface terms (see Eq. (42)).

We are now almost ready to write the action with the new canonical variables. It is now necessary to determine the new Lagrange multipliers. In order to write the new constraints with the new Lagrange multipliers, we can use the identity given by the space derivative of M,

$$M' = -\Lambda^{-1}\left(R'H + \frac{1}{2}P_\Lambda H_r\right)\,. \qquad (74)$$

Solving for H and making use of the inverse transformations of Λ and P_Λ, in Eqs. (59) and (60), we get

$$\begin{aligned} H &= -\frac{M'F^{-1}\mathrm{R}' + FP_M P_{\mathrm{R}}}{\left(F^{-1}(\mathrm{R}')^2 - FP_M^2\right)^{\frac{1}{2}}}\,, && (75)\\ H_r &= P_M M' + P_{\mathrm{R}}\mathrm{R}'\,. && (76) \end{aligned}$$

Following Kuchař [11], the new set of constraints, totally equivalent to the old set $H(t,r) = 0$ and $H_r(t,r) = 0$ outside the horizon points, is $M'(t,r) = 0$ and $P_{\mathrm{R}}(t,r) = 0$. By continuity, this also applies on the horizon, where $F(t,r) = 0$. So we can say that the equivalence is valid everywhere. The new Hamiltonian, the total sum of the constraints, can now be written as

$$NH + N^r H_r = N^M M' + N^{\mathrm{R}} P_{\mathrm{R}}\,. \qquad (77)$$

In order to determine the new Lagrange multipliers, one has to write the left hand side of the previous equation, Eq. (77), and replace the constraints on that side by their expressions as functions of the new canonical coordinates, spelt out in Eqs. (75)-(76). After manipulation, one gets

$$\begin{aligned} N^M &= -\frac{NF^{-1}R'}{\left(F^{-1}(\mathrm{R}')^2 - FP_M^2\right)^{\frac{1}{2}}} + N^r P_M\,, && (78)\\ N^{\mathrm{R}} &= -\frac{NFP_M}{\left(F^{-1}(\mathrm{R}')^2 - FP_M^2\right)^{\frac{1}{2}}} + N^r R'\,. && (79) \end{aligned}$$

Using the inverse transformations Eqs. (59)-(60), and the identity $R = \mathrm{R}$, we can write the new multipliers as functions of the old variables

$$\begin{aligned} N^M &= -NF^{-1}R'\Lambda^{-1} + \frac{1}{2}N^r F^{-1}\Lambda P_\Lambda\,, && (80)\\ N^{\mathrm{R}} &= -\frac{1}{2}NP_\Lambda + N^r R'\,, && (81) \end{aligned}$$

allowing us determine its asymptotic conditions from the original conditions given above. These transformations are non-singular for $r > 0$. As before, for $r \to 0$,

$$\begin{aligned} N^M(t,r) &= -\frac{1}{2}N_1(t)\Lambda_0 R_2^{-1} + O(r^2)\,, && (82)\\ N^{\mathrm{R}}(t,r) &= O(r^4)\,, && (83) \end{aligned}$$

and for $r \to \infty$ we have

$$\begin{aligned} N^M(t,r) &= -\tilde{N}_+(t) + O^\infty(r^{-4})\,, && (84)\\ N^{\mathrm{R}}(t,r) &= O^\infty(r^{-1})\,. && (85) \end{aligned}$$

The conditions (82)-(85) show that the transformations in Eqs. (80)-(81) are satisfactory in the case of $r \to \infty$, but not for $r \to 0$. This is due to fact that in order to fix the Lagrange multipliers for $r \to \infty$, as we are free to do, we fix $\tilde{N}_+(t)$, which we already do when adding the surface term

$$-\int dt\, \tilde{N}_+ M_+ \qquad (86)$$

to the action, in order to obtain the equations of motion in the bulk, without surface terms. However, at $r=0$, we see that fixing the multiplier N^M to values independent of the canonical variables is not the same as fixing $N_1\Lambda_0^{-1}$ to values independent of the canonical variables. We need to rewrite the multiplier N^M for the asymptotic regime $r\rightarrow 0$ without affecting its behavior for $r\rightarrow\infty$. In order to proceed we have to make one assumption, which is that the expression given in asymptotic condition of $M(t,r)$, as $r\rightarrow 0$, for the term of order zero, $M_0\equiv 2\lambda^2 R_0(t)^2$, defines R_0 as a function of M_0, and R_0 is the horizon radius function, $R_0\equiv R_{\mathrm{h}}(M_0)$. Also, we assume that $M_0>0$. With these assumptions, we are working in the domain of the classical solutions. We can immediately obtain that the variation of R_0 is given in relation to the variation of M_0 as

$$\delta R_0 = (2\lambda)^{-2} R_0^{-1}\delta M_0\,, \tag{87}$$

where, as defined above, $a^2 = l^{-2} = 2\lambda^2$. This expression will be used when we derive the equations of motion from the new action. We now define the new multiplier $\tilde{N}^M$ as

$$\tilde{N}^M = -N^M\left[(1-g) + gR_0\left(l^{-2}R_0^2\right)^{-1}\right]^{-1}\,, \tag{88}$$

where $g(r) = 1 + O(r^2)$ for $r\rightarrow 0$ and $g(r) = O^\infty(r^{-5})$ for $r\rightarrow\infty$. This new multiplier, function of the old multiplier N^M, has as its properties for $r\rightarrow\infty$

$$\tilde{N}^M(t,r) = \tilde{N}_+(t) + O^\infty(r^{-4})\,, \tag{89}$$

and as its properties for $r\rightarrow 0$

$$\tilde{N}^M(t,r) = \tilde{N}_0^M(t) + O(r^2)\,, \tag{90}$$

where $\tilde{N}_0^M$ is given by

$$\tilde{N}_0^M = \lambda^2 N_1 R_0 R_2^{-1}\Lambda_0\,. \tag{91}$$

When the constraint $M'=0$ holds, the last expression is

$$\tilde{N}_0^M = N_1\Lambda_0^{-1}\,. \tag{92}$$

With this new constraint $\tilde{N}^M$, fixing $N_1\Lambda_0^{-1}$ at $r=0$ or fixing $\tilde{N}^M$ is equivalent, there being no problems with N^{R}, which is left as determined in Eq. (79).

The new action is now written as the sum of S_Σ, the bulk action, and $S_{\partial\Sigma}$, the surface action,

$$\begin{aligned} S\left[M,\mathrm{R},P_M,P_{\mathrm{R}};\tilde{N}^M,N^{\mathrm{R}}\right] \;=\; & \int dt\int_0^\infty dr\,\left(P_M\dot{M} + P_{\mathrm{R}}\dot{\mathrm{R}} - N^{\mathrm{R}}P_{\mathrm{R}} + \tilde{N}^M\left[(1-g)+gR_0\left(l^{-2}R_0^2\right)^{-1}\right]M'\right) + \\ & \int dt\,\left(2R_0\tilde{N}_0^M - \tilde{N}_+M_+\right)\,. \end{aligned} \tag{93}$$

The new equations of motion are now

$$\dot{M} \;=\; 0\,, \tag{94}$$

$$\dot{\mathrm{R}} \;=\; N^{\mathrm{R}}\,, \tag{95}$$

$$\dot{P}_M \;=\; (N^M)'\,, \tag{96}$$

$$\dot{P}_{\mathrm{R}} \;=\; 0\,, \tag{97}$$

$$M' \;=\; 0\,, \tag{98}$$

$$P_{\mathrm{R}} \;=\; 0\,, \tag{99}$$

where we understood N^M to be a function of the new constraint, defined through Eq. (88). The resulting boundary terms of the variation of this new action, Eq. (93), are, first, terms proportional to δM and $\delta\mathrm{R}$ on the initial and final hypersurfaces, and, second, $\int dt\,\left(2R_0\delta\tilde{N}_0^M - M_+\delta\tilde{N}_+\right)$. Here we have used the expression in Eq. (87). The action in Eq. (93) yields the equations of motion, Eqs. (94)-(99), provided that we fix the initial and final values of the new canonical variables and that we also fix the values of $\tilde{N}_0^M$ and of $\tilde{N}_+$. Thanks to the redefinition of the Lagrange multiplier, from N^M to $\tilde{N}^M$, the fixation of those quantities, $\tilde{N}_0^M$ and $\tilde{N}_+$, has the same meaning it had before the canonical transformations and the redefinition of N^M. This same meaning is guaranteed through the use of our gauge freedom to choose the multipliers, and at the same time not fixing the boundary variations independently of the choice of Lagrange multipliers, which in turn allow us to have a well defined variational principle for the action.

D. Hamiltonian reduction

We now solve the constraints in order to reduce to the true dynamical degrees of freedom. The equations of motion (94)-(99) allow us to write M as an independent function of space, r,

$$M(t,r) = \mathbf{m}(t)\,. \tag{100}$$

The reduced action, with the constraints taken into account, is then

$$S\left[\mathbf{m}, \mathbf{p_m}; \tilde{N}_0^M, \tilde{N}_+\right] = \int dt\, \mathbf{p_m}\dot{\mathbf{m}} - \mathbf{h}\,, \tag{101}$$

where

$$\mathbf{p_m} = \int_0^\infty dr\, P_M\,, \tag{102}$$

and the reduced Hamiltonian, $\mathbf{h}$, is now written as

$$\mathbf{h}(\mathbf{m};t) = -2R_{\mathrm{h}}\tilde{N}_0^M + \tilde{N}_+\mathbf{m}\,, \tag{103}$$

with R_{h} being the horizon radius. We also have that $\mathbf{m} > 0$, according to the assumptions made in the previous subsection. Thanks to the functions $\tilde{N}_0^M(t)$ and $\tilde{N}_+(t)$ the Hamiltonian $\mathbf{h}$ is an explicitly time dependent function. The variational principle associated with the reduced action, Eq. (101), will fix the value of $\mathbf{m}$ on the initial and final hypersurfaces, or in the spirit of the classical analytical mechanics, the Hamiltonian principle fixes the initial and final values of the canonical coordinate. The equations of motion are

$$\dot{\mathbf{m}} = 0\,, \tag{104}$$

$$\dot{\mathbf{p}}_{\mathbf{m}} = 2\tilde{N}_0^M(4\lambda^2 R_{\mathrm{h}})^{-1} - \tilde{N}_+\,. \tag{105}$$

The equation of motion for $\mathbf{m}$, Eq. (104), is understood as saying that $\mathbf{m}$ is, on a classical solution, equal to the mass parameter M of the solution, Eq. (16). In order to interpret the other equation of motion, Eq. (105), we have to recall that from Eq. (57) one has $P_M = -T'$, where T is the Killing time. This, together with the definition of $\mathbf{p_m}$, given in Eq. (102), yields

$$\mathbf{p_m} = T_0 - T_+\,, \tag{106}$$

where T_0 is the value of the Killing time at the left end of the hypersurface of a certain t, and T_+ is the Killing time at spatial infinity, the right end of the same hypersurface of t. As the hypersurface evolves in the spacetime of the black hole solution, the right hand side of Eq. (105) is equal to $\dot{T}_0 - \dot{T}_+$.

E. Quantum theory and partition function

The next step is to quantize the reduced Hamiltonian theory, by building the time evolution operator quantum mechanically and then obtaining a partition function through the analytic continuation of the same operator [12]-[16]. The variable $\mathbf{m}$ is regarded here as a configuration variable. This variable satisfies the inequality $\mathbf{m} > 0$. The wave functions will be of the form $\psi(\mathbf{m})$, with the inner product given by

$$(\psi, \chi) = \int_A \mu d\mathbf{m}\, \bar{\psi}\chi\,, \tag{107}$$

where A is the domain of integration defined by $\mathbf{m} > 0$ and $\mu(\mathbf{m})$ is a smooth and positive weight factor for the integration measure. It is assumed that μ is a slow varying function, otherwise arbitrary. We are thus working in the Hilbert space defined as $\mathscr{H} := L^2(A; \mu d\mathbf{m})$.

The Hamiltonian operator, written as $\hat{\mathbf{h}}(t)$, acts through pointwise multiplication by the function $\mathbf{h}(\mathbf{m};t)$, which on a function of our working Hilbert space reads

$$\hat{\mathbf{h}}(t)\psi(\mathbf{m}) = \mathbf{h}(\mathbf{m};t)\psi(\mathbf{m})\,. \tag{108}$$

This Hamiltonian operator is an unbounded essentially self-adjoint operator. The corresponding time evolution operator in the same Hilbert space, which is unitary due to the fact that the Hamiltonian operator is self-adjoint, is

$$\hat{K}(t_2;t_1) = \exp\left[-i\int_{t_1}^{t_2} dt'\,\hat{\mathbf{h}}(t')\right]\,. \tag{109}$$

This operator acts also by pointwise multiplication in the Hilbert space. We now define

$$\mathcal{T} \;:=\; \int_{t_1}^{t_2} dt\,\tilde{N}_+(t)\,, \tag{110}$$

$$\Theta \;:=\; \int_{t_1}^{t_2} dt\,\tilde{N}_0^M(t)\,. \tag{111}$$

Using (103), (109), (110), and (111), we write the function K, which is in fact the action of the operator in the Hilbert space, as

$$K\left(\mathbf{m};\mathcal{T},\Theta\right) = \exp\left[-i\mathbf{m}\mathcal{T} + 2\,i\,R_{\rm h}\Theta\right]\,. \tag{112}$$

This expression indicates that $\hat{K}(t_2;t_1)$ depends on t_1 and t_2 only through the functions $\mathcal{T}$ and Θ. Thus, the operator corresponding to the function K can now be written as $\hat{K}(\mathcal{T};\Theta)$. The composition law in time $\hat{K}(t_3;t_2)\hat{K}(t_2;t_1) = \hat{K}(t_3;t_1)$ can be regarded as a sum of the parameters $\mathcal{T}$ and Θ, inside the operator $\hat{K}(\mathcal{T};\Theta)$. These parameters are evolutions parameters defined by the boundary conditions, i.e., $\mathcal{T}$ is the Killing time elapsed at right spatial infinity and Θ is the boost parameter elapsed at the bifurcation circle.

F. Thermodynamics

We can now build the partition function for this system. The path to follow is to continue the operator to imaginary time and take the trace over a complete orthogonal basis. Our classical thermodynamic situation consists of a three-dimensional spherically symmetric black hole, asymptotically anti-de Sitter, in thermal equilibrium with a bath of Hawking radiation. Ignoring back reaction from the radiation, the geometry is described by the solutions in Eq. (16). Thus, we consider a thermodynamic ensemble in which the temperature, or more appropriately here, the inverse temperature β is fixed. This characterizes a canonical ensemble, and the partition function $\mathcal{Z}(\beta)$ arises naturally in such an ensemble. To analytically continue the Lorentzian solution we put $\mathcal{T} = -i\beta$, and $\Theta - 2\pi i$, this latter choice based on the regularity of the classical Euclidean solution. We arrive then at the following expression for the partition function

$$\mathcal{Z}(\beta) = \mathrm{Tr}\left[\hat{K}(-i\beta,-2\pi i)\right]\,. \tag{113}$$

From Eq. (112) this is realized as

$$\mathcal{Z}(\beta) = \int_0^\infty \mu\,d\mathbf{m}\,\exp\left[-\beta\mathbf{m} + 4\pi R_{\rm h}\right]\langle\mathbf{m}|\mathbf{m}\rangle\,. \tag{114}$$

Since $\langle\mathbf{m}|\mathbf{m}\rangle$ is equal to $\delta(0)$, one has to regularize Eq. (114). Following the procedure developed in the Louko-Whiting approach [12], this means regularizing and normalizing the operator $\hat{K}$ beforehand. This leads to

$$\mathcal{Z}_{\rm ren}(\beta) = \mathcal{N}\int_A \mu\,d\mathbf{m}\,\exp\left[-\beta\mathbf{m} + 4\pi R_{\rm h}\right]\,, \tag{115}$$

where $\mathcal{N}$ is a normalization factor and A is the domain of integration. Provided the weight factor μ is slowly varying compared to the exponential in Eq. (115), and using the fact that the horizon radius $R_{\rm h}$ is a function of $\mathbf{m}$, the integral in Eq. (115) is convergent. Thermodynamically, $\mathbf{m}$ is analogous to the energy of the system. Changing integration variables, from $\mathbf{m}$ to $R_{\rm h}$, where

$$\mathbf{m} = 2\lambda^2 R_{\rm h}^2\,, \tag{116}$$

the integral Eq. (115) becomes

$$\mathcal{Z}_{\rm ren}(\beta) = \mathcal{N}\int_{A'} \widetilde{\mu}\,dR_{\rm h}\,\exp(-I_*)\,, \tag{117}$$

where A' is new the domain of integration after changing variables, and the function $I_*(R_{\rm h})$, a kind of an effective action (see [4]), is written as

$$I_*(R_{\rm h}) := 2\beta\,\lambda^2 R_{\rm h}^2 - 4\pi R_{\rm h}\,. \tag{118}$$

The new domain of integration, A', is defined by the inequality $R_{\rm h} \geq 0$. The new weight factor $\widetilde{\mu}$ includes the Jacobian of the transformation, which amounts to $\partial \mathbf{m}/\partial R_{\rm h}$. Since the weight factor $\widetilde{\mu}$ is slowly varying, we can estimate the integral of $\mathcal{Z}_{\rm ren}(\beta)$ by the saddle point approximation. For that we have to calculate the critical points, i.e., we have to find the values of $R_{\rm h}$ for which the first derivative of $I_*(R_{\rm h})$ with respect to $R_{\rm h}$ is zero. It happens for only one value of the domain

$$R_{\rm h}^+ \;=\; \frac{\pi}{\beta\,\lambda^2}\,. \tag{119}$$

In order to find out the nature of the extremum we evaluate the second derivative at the extremum. One finds

$$\left.\frac{\partial^2 I_*}{\partial R_{\rm h}^2}\right|_{R_{\rm h}^+} \;=\; 4\beta\lambda^2\,. \tag{120}$$

As our domain starts at $R_{\rm h} = 0$, we have that the extremum located at $R_{\rm h} = R_{\rm h}^+$ is a minimum. Evaluating the action I_* at $R_{\rm h}^+$ one obtains

$$I_*(R_{\rm h}^+) = -\frac{2\pi^2}{\beta\lambda^2} < 0\,. \tag{121}$$

By Taylor expanding the action in the exponential of the integral, Eq. (117),

$$I_*(R_{\rm h}) = I_*(R_{\rm h}^+) + \left.\frac{\partial I_*}{\partial R_{\rm h}}\right|_{R_{\rm h}^+} R_{\rm h} + \frac{1}{2}\left.\frac{\partial^2 I_*}{\partial R_{\rm h}^2}\right|_{R_{\rm h}^+} (R_{\rm h})^2 + O((R_{\rm h})^3)\,, \tag{122}$$

we can separate the terms in such a way that we obtain the following expression for the renormalized partition function

$$\mathcal{Z}_{\rm ren}(\beta) = \exp\left[-I_*(R_{\rm h}^+)\right] \mathcal{N} \int_{A'} \widetilde{\mu}\, dR_{\rm h} \exp\left[2\beta\lambda^2 R_{\rm h}^2\right]\,. \tag{123}$$

The Taylor expansion was up to second order, and evaluated at the critical point $R_{\rm h}^+$, which makes the first order term of the expansion of $I_*(R_{\rm h})$ disappear. The term which can be put outside the integral is the zero order term, which is the value of $I_*(R_{\rm h})$ at the extremum $R_{\rm h}^+$. The term left inside the exponential, $2\beta\lambda^2 R_{\rm h}^2$, is minus the second order term in the Taylor expansion, where all the higher orders have been ignored, as this is a good approximation, provided the weight factor is slowly varying. Finally, we may write the renormalized partition function as

$$\mathcal{Z}_{\rm ren}(\beta) = {\rm P}\exp\left(\frac{2\pi^2}{\beta\lambda^2}\right)\,, \tag{124}$$

where P is given by

$${\rm P} = \mathcal{N} \int_{A'} \widetilde{\mu}\, dR_{\rm h} \exp\left(2\beta\lambda^2 R_{\rm h}^2\right)\,. \tag{125}$$

This P is a slowly varying prefactor and this approximation is better as we move to higher values of $|I_*(R_{\rm h}^+)|$. In the domain of integration the dominating contribution comes from the vicinity of $R_{\rm h} = R_{\rm h}^+$. Leaving the explicit dependence of the partition function on the $R_{\rm h}^+$ we write the logarithm of $\mathcal{Z}_{\rm ren}(\beta)$ as

$$\ln(\mathcal{Z}_{\rm ren}(\beta)) = \ln {\rm P} - 2\beta\,\lambda^2 (R_{\rm h}^+)^2 + 4\pi R_{\rm h}^+\,. \tag{126}$$

By ignoring the prefactor's logarithm, which closer to $R_{\rm h}^+$ is less relevant, we are able to determine the value of $\mathbf{m}$ at the critical point, where we find that it corresponds to the value of the mass of the classical solutions of the black holes given in Eq. (16). Thus, when the critical point dominates the partition function, we have that the mean energy $\langle E\rangle$ is given by

$$\langle E\rangle = -\frac{\partial}{\partial\beta}\ln \mathcal{Z}_{\rm ren}(\beta) \approx 2\lambda^2 (R_{\rm h}^{+})^2 = \mathbf{m}^{+}\,, \tag{127}$$

where $\mathbf{m}^{+}$ is obtained from Eq. (116) evaluated at $R_{\rm h}^{+}$. Using Eq. (119) and Eq. (116), we obtain the temperature of the black hole $\mathbf{T}\equiv\beta^{-1}$

$$\mathbf{T} = \left(\frac{\lambda^2\mathbf{m}^{+}}{2\pi^2}\right)^{\frac{1}{2}}\,, \tag{128}$$

where $\mathbf{m}^{+}$ is the function in (116) evaluated at $R_{\rm h}^{+}$. Inverting Eq. (128), and writing the inverse temperature as β, we write the function $\mathbf{m}^{+}(\beta)$

$$\mathbf{m}^{+}(\beta) = \frac{2\pi^2}{\beta^2\lambda^2}\,. \tag{129}$$

We see that $\partial\mathbf{m}^{+}/\partial\beta < 0$, which through $C = -\beta^2(\partial\langle E\rangle/\partial\beta)$ tells us that the system is thermodynamically stable. The entropy is given by

$$S = \left(1-\beta\frac{\partial}{\partial\beta}\right)(\ln \mathcal{Z}_{\rm ren}(\beta)) \approx 4\pi R_{\rm h}^{+}\,. \tag{130}$$

This is the entropy of the BTZ black hole (see [20] and also [26, 27, 28]).

IV. HAMILTONIAN THERMODYNAMICS OF THE GENERAL RELATIVISTIC CYLINDRICAL DIMENSIONALLY REDUCED BLACK HOLE ($\omega = 0$)

A. The metric

For $\omega = 0$, the corresponding three-dimensional Brans-Dicke theory is obtained from the cylindrical dimensionally reduced black hole of four-dimensional general relativity [24, 25]. Then general metric in Eq. (2) and the ϕ field in Eq. (6), reduce to the following

$$\begin{aligned} ds^2 &= -\left[(a\,R)^2 - \frac{M}{2(a\,R)}\right]dT^2 + \frac{dR^2}{(a\,R)^2 - \frac{M}{2(a\,R)}} + R^2\,d\varphi^2\,, && (131)\\ e^{-2\phi} &= a\,R\,, && (132) \end{aligned}$$

with $M = 2b$ and $a = \sqrt{\frac{2}{3}}|\lambda|$. Unlike the BTZ solution, this solution (131)-(132) has a metric function whose mass term depends on R. This behavior is similar to the Schwarzschild black hole metric function [24]. In Fig. 2 we have plotted the Carter-Penrose diagram of the black hole solution for $\omega = 0$, where again $R_{\rm h}$ is the horizon radius, given by the larger positive real root of $R_{\rm h}^3 = \frac{1}{2\,a^3}M$, $R = 0$ is the radius of the curvature singularity, and $R = \infty$ is the spatial infinity. The roman numerals mean the same as in Fig. 1.

B. Canonical formalism

The action with $\omega = 0$ becomes, excluding surface terms,

$$\begin{aligned} S[\Lambda,\,R,\,\dot\Lambda,\,\dot R;\,N,\,N^r] = \int dt\int_0^\infty dr\,\alpha\Big\{&\lambda^2\Lambda N R^2 - N^{-1}R\dot R\dot\Lambda - \frac{1}{2}N^{-1}\Lambda\dot R^2 + N^{-1}R\dot R(\Lambda N^r)'\\ &+N^{-1}N^r R R'\dot\Lambda - N^{-1}N^r\Lambda R R'(N^r)' - N^{-1}(N^r)^2 R R'\Lambda' + N^{-1}N^r\Lambda\dot R R'\\ &-\frac{1}{2}N^{-1}(N^r)^2\Lambda(R')^2 - (\Lambda^{-1})' R R' N - \frac{1}{2}\Lambda^{-1}(R')^2 N - \Lambda^{-1}RR''N\Big\}\,, \end{aligned} \tag{133}$$



FIG. 2: The Carter-Penrose Diagram for the $\omega = 0$ case.

where $\dot{}$ means derivative with respect to time t and $'$ is the derivative with respect to r, and where all the explicit functional dependences are omitted. Depending on the situation we use four different letters containing the same information but with slightly different numerical values. Thus, a, λ, l and α are related by $a^2 = \frac{2}{3}\lambda^2 = l^{-2}$, and $\alpha = 4a$, where l is, as usual, the AdS length. From this action one determines the canonical momenta, conjugate to Λ and R respectively

$$\begin{aligned} P_\Lambda &= -\alpha N^{-1} R \left\{ \dot{R} - R' N^r \right\} \,, && (134) \\ P_R &= -\alpha N^{-1} \left\{ R[\dot{\Lambda} - (\Lambda N^r)'] + \Lambda[\dot{R} - N^r R'] \right\} \,. && (135) \end{aligned}$$

By performing a Legendre transformation, we obtain

$$\begin{aligned} \mathcal{H} &= N \left\{ -\alpha^{-1} R^{-1} P_\Lambda P_R + \frac{1}{2}\alpha^{-1}\Lambda R^{-2} P_\Lambda^2 + \alpha \Lambda^{-1} R R'' - \alpha \Lambda^{-2} R R' \Lambda' + \frac{1}{2}\Lambda^{-1}(R')^2 - \lambda^2 \Lambda R^2 \right\} \\ &\quad + N^r \left\{ P_R R' - P_\Lambda' \Lambda \right\} \equiv N H + N^r H_r \,. && (136) \end{aligned}$$

The action in Hamiltonian form is then

$$S[\Lambda,\, R,\, P_\Lambda,\, P_R;\, N,\, N^r] = \int dt \int_0^\infty dr \left\{ P_\Lambda \dot{\Lambda} + P_R \dot{R} - N H - N^r H_r \right\} \,. \qquad (137)$$

The equations of motion are

$$\begin{aligned} \dot{\Lambda} &= -N\alpha^{-1} R^{-1} P_R + N\alpha^{-1}\Lambda R^{-2} P_\Lambda + N^r \Lambda && (138) \\ \dot{R} &= -N\alpha^{-1} P_\Lambda R^{-1} + N^r R' && (139) \\ \dot{P_R} &= -N\alpha^{-1} P_\Lambda P_R R^{-2} + N\alpha^{-1}\Lambda P_\Lambda^2 R^{-3} - \left((N\alpha)' \Lambda^{-1} R \right)' - (N\alpha)(\Lambda^{-1} R')' + 2N\alpha\lambda^2 \Lambda R + (N^r P_R)' && (140) \\ \dot{P_\Lambda} &= -\frac{1}{2} N\alpha^{-1} R^{-2} P_\Lambda^2 - (N\alpha)' R R' \Lambda^{-2} - \frac{1}{2} N\alpha (R')^2 \Lambda^{-2} + N\alpha\lambda^2 R^2 + N^r P_\Lambda' \,. && (141) \end{aligned}$$

In order to have a well defined variational principle, we need to eliminate the surface terms of the original bulk action, which render the original action itself ill defined for a correct determination of the equations of motion through variational methods. Through the choice of added surface terms one can achieve this elimination. The action (137) has the following extra surface terms, after variation

$$\text{Surface terms} = \alpha \left(-N\Lambda^{-1} R \delta R' + N' \Lambda^{-1} R \delta R - N^r P_R \delta R + N^r \Lambda \delta P_\Lambda + N R R' \Lambda^{-2} \delta\Lambda \right)\Big|_0^\infty \,. \qquad (142)$$

In order to evaluate this expression, we need to know the asymptotic conditions of each of the functions of (t, r).

Starting with the limit $r \to 0$ we assume

$$\begin{aligned} \Lambda(t,r) &= \Lambda_0(t) + O(r^2) \,, && (143) \\ R(t,r) &= R_0(t) + R_2(t) r^2 + O(r^4) \,, && (144) \\ P_\Lambda(t,r) &= O(r^3) && (145) \\ P_R(t,r) &= O(r) \,, && (146) \\ N(t,r) &= N_1(t) r + O(r^3) && (147) \\ N^r(t,r) &= N_1^r(t) r + O(r^3) \,. && (148) \end{aligned}$$

With these conditions, we have for the surface terms at $r=0$,

$$\text{Surface terms}\big|_{r=0} = -\alpha N_1 R_0 \Lambda_0^{-1} \delta R_0 \,. \tag{149}$$

In the same way, for $r \to \infty$,

$$\begin{aligned}
\Lambda(t,r) &= l r^{-1} + l^3 \eta(t) r^{-4} + O^\infty(r^{-5})\,, && (150)\\
R(t,r) &= r + l^2 \rho(t) r^{-2} + O^\infty(r^{-3})\,, && (151)\\
P_\Lambda(t,r) &= O^\infty(r^{-2}) && (152)\\
P_R(t,r) &= O^\infty(r^{-4})\,, && (153)\\
N(t,r) &= R(t,r)' \Lambda(t,r)^{-1} (\tilde{N}_+(t) + O^\infty(r^{-5}))\,, && (154)\\
N^r(t,r) &= O^\infty(r^{-2})\,. && (155)
\end{aligned}$$

These conditions imply for the surface terms in the limit $r \to \infty$,

$$\text{Surface terms}\big|_{r\to\infty} = \alpha \delta(M_+) \tilde{N}_+ \,, \tag{156}$$

where $M_+(t) = \alpha(\eta(t) + 3\rho(t))$. So, the surface term to be added to (137) is

$$S_{\partial\Sigma}\left[\Lambda, R; N\right] = \int dt \left(\frac{1}{2} \alpha R_0^2 N_1 \Lambda_0^{-1} - \tilde{N}_+ M_+ \right) . \tag{157}$$

What is left after varying this last surface term and adding it to the varied initial action (see Eq. (137)) is

$$\int dt \left(\frac{1}{2} \alpha R_0^2 \delta(N_1 \Lambda_0^{-1}) - \delta\tilde{N}_+ M_+ \right) . \tag{158}$$

We choose to fix $N_1 \Lambda_0^{-1}$ on the horizon and $\tilde{N}_+$ at infinity, which makes the surface variation (158) disappear.

C. Reconstruction, canonical transformation, and action

In order to reconstruct the mass and the time from the canonical data, which amounts to making a canonical transformation, we have to rewrite the general form of the solutions in Eqs. (131)-(132). We again follow Kuchař [11] for the reconstruction. We concentrate our analysis in the right static region of the Carter-Penrose diagram. In the right static region we define F as

$$F(R(t,r)) = (aR(t,r))^2 - \frac{b}{aR(t,r)}\,, \tag{159}$$

and make the following substitutions

$$T = T(t,r)\,, \qquad\qquad R = R(t,r)\,, \tag{160}$$

in the solutions above, Eqs. (131)-(132), getting

$$ds^2 \;=\; -(F\dot{T}^2 - F^{-1}\dot{R}^2)\,dt^2 \;+\; 2(-FT'\dot{T} + F^{-1}R'\dot{R})\,dtdr \;+\; (-F(T')^2 + F^{-1}\dot{R}^2)\,dr^2 \;+\; R^2 d\varphi^2\,. \tag{161}$$

This introduces the ADM foliation directly into the solutions. Comparing it with the ADM metric (14), written in another form as

$$ds^2 = -(N^2 - \Lambda^2 (N^r)^2)\,dt^2 \,+\, 2\Lambda^2 N^r \,dtdr \,+\, \Lambda^2 dr^2 \,+\, R^2\, d\varphi^2\,, \tag{162}$$

we can write a set of three equations

$$\begin{aligned}
\Lambda^2 &= -F(T')^2 + F^{-1}(R')^2\,, && (163)\\
\Lambda^2 N^r &= -FT'\dot{T} + F^{-1}R'\dot{R}\,, && (164)\\
N^2 - \Lambda^2 (N^r)^2 &= F\dot{T}^2 - F^{-1}\dot{R}^2\,. && (165)
\end{aligned}$$

The first two equations, Eq. (163) and Eq. (164), give

$$N^r = \frac{-FT'\dot{T} + F^{-1}R'\dot{R}}{-F(T')^2 + F^{-1}(R')^2}\,. \tag{166}$$

This one solution, together with Eq. (163), give

$$N = \frac{R'\dot{T} - T'\dot{R}}{\sqrt{-F(T')^2 + F^{-1}(R')^2}}\,. \tag{167}$$

One can show that $N(t,r)$ is positive. Next, putting Eq. (166) and Eq. (167) into the definition of the conjugate momentum of the canonical coordinate Λ, given in Eq. (134), one finds the spatial derivative of $T(t,r)$ as a function of the canonical coordinates, i.e.,

$$-T' = \alpha^{-1}R^{-1}F^{-1}\Lambda P_\Lambda\,. \tag{168}$$

Later we will see that $-T' = P_M$, as it will be conjugate to a new canonical coordinate M. Following this procedure to the end, we may then find the form of the new coordinate $M(t,r)$, also as a function of t and r. First, we need to know the form of F as a function of the canonical pair Λ, R. For that, we replace back into Eq. (163) the definition of T', giving

$$F = \left(\frac{R'}{\Lambda}\right)^2 - \left(\frac{P_\Lambda}{\alpha R}\right)^2\,. \tag{169}$$

Equating this form of F with Eq. (159), we obtain

$$M = \frac{1}{2}\alpha R\left(\frac{\alpha^2}{16}R^2 - F\right)\,, \tag{170}$$

where F is given in Eq. (169). We thus have found the form of the new canonical coordinate, M. It is now a straightforward calculation to determine the Poisson bracket of this variable with $P_M = -T'$ and see that they are conjugate, thus making Eq. (168) the conjugate momentum of M, i.e.,

$$P_M = \alpha^{-1}R^{-1}F^{-1}\Lambda P_\Lambda\,. \tag{171}$$

It is now necessary to find out the other canonical variable which commutes with M and P_M and which guarantees, plus its conjugate momentum, that the transformation from Λ, R to M and the new variable is canonical. Immediately is it seen that R commutes with M and P_M. It is then a candidate. It remains to be seen whether P_R also commutes with M and P_M. As with R, it is straightforward to see that P_R does not commute with M and P_M, as these contain powers of R in their definitions, and $\{R(t,r),\, P_R(t,r^*)\} = \delta(r - r^*)$. So rename the new canonical variable R as $R = \mathrm{R}$. We have to find a new conjugate momentum to R which also commutes with M and P_M, making the transformation from $\{\Lambda,\, R;\, P_\Lambda,\, P_R\} \rightarrow \{M,\, \mathrm{R};\, P_M,\, P_{\mathrm{R}}\}$ a canonical one. The way to proceed is to look at the constraint H_r, which is called in this formalism the super-momentum. This is the constraint which generates spatial diffeomorphisms in all variables. Its form, in the initial canonical coordinates, is $H_r = P_R\,R' - \Lambda\,P_\Lambda'$. In this formulation, Λ is a spatial density and R is a spatial scalar. As all new variables, M and R, are spatial scalars, the generator of spatial diffeomorphisms is written as $H_r = P_{\mathrm{R}}\mathrm{R}' + P_M M'$, regardless of the particular form of the canonical coordinate transformation. It is thus equating these two expressions of the super-momentum H_r, with M and P_M written as functions of Λ, R and their respective momenta, that gives us the equation for the new P_{R}. This results in

$$\begin{aligned} P_{\mathrm{R}} \;=\;& P_R - \frac{3\alpha^2}{32}F^{-1}\Lambda P_\Lambda R - \frac{1}{2}R^{-1}\Lambda P_\Lambda + F^{-1}P_\Lambda R''\Lambda^{-1} - F^{-1}\Lambda^{-2}P_\Lambda\Lambda' R' + (R')^2F^{-1}\Lambda^{-1}P_\Lambda R^{-1} \\ & -F^{-1}\Lambda^{-1}P_\Lambda' R'\,. \end{aligned} \tag{172}$$

We have now all the canonical variables of the new set determined. For completeness and future use, we write the inverse transformation for Λ and P_Λ,

$$\Lambda \;=\; \left((\mathrm{R}')^2F^{-1} - P_M^2 F\right)^{\frac{1}{2}}\,, \tag{173}$$

$$P_\Lambda \;=\; \alpha \mathrm{R} F P_M\left((\mathrm{R}')^2F^{-1} - P_M^2 F\right)^{-\frac{1}{2}}\,. \tag{174}$$

In summary, the canonical transformations are

$$\begin{aligned}
R &= \mathrm{R}\,,\\
M &= \frac{1}{2}\alpha R\left(\frac{\alpha^2}{16}R^2-F\right)\,,\\
P_{\mathrm{R}} &= P_R-\frac{3\alpha^2}{32}F^{-1}\Lambda P_\Lambda R-\frac{1}{2}R^{-1}\Lambda P_\Lambda+F^{-1}P_\Lambda R''\Lambda^{-1}-F^{-1}\Lambda^{-2}P_\Lambda\Lambda' R'+(R')^2F^{-1}\Lambda^{-1}P_\Lambda R^{-1}\\
&\quad -F^{-1}\Lambda^{-1}P_\Lambda' R'\,,\\
P_M &= \alpha^{-1}R^{-1}F^{-1}\Lambda P_\Lambda\,. \qquad (175)
\end{aligned}$$

In order to prove that the set of equalities in expression (175) is canonical we start with the equality

$$\begin{aligned}
P_\Lambda\delta\Lambda+P_R\delta R-P_M\delta M-P_{\mathrm{R}}\delta\mathrm{R} &= \left(\frac{1}{2}\alpha R\delta R\ln\left|\frac{\alpha RR'+\Lambda P_\Lambda}{\alpha RR'-\Lambda P_\Lambda}\right|\right)'+\\
&\quad+\delta\left(\Lambda P_\Lambda+\frac{1}{2}\alpha RR'\ln\left|\frac{\alpha RR'-\Lambda P_\Lambda}{\alpha RR'+\Lambda P_\Lambda}\right|\right)\,. \qquad (176)
\end{aligned}$$

We now integrate expression (176) in r, in the interval from $r=0$ to $r=\infty$. The first term on the right hand side of Eq. (176) vanishes due to the falloff conditions (see Eqs. (143)-(148) and Eqs. (150)-(155)). We then obtain the following expression

$$\int_0^\infty dr\ \left(P_\Lambda\delta\Lambda+P_R\delta R\right)-\int_0^\infty dr\ \left(P_M\delta M+P_{\mathrm{R}}\delta\mathrm{R}\right) = \delta\omega\left[\Lambda,\,R,\,P_\Lambda\right]\,, \qquad (177)$$

where $\delta\omega\left[\Lambda,\,R,\,P_\Lambda\right]$ is a well defined functional, which is also an exact form. This equality shows that the difference between the Liouville form of $\{R,\,\Lambda;\,P_R,\,P_\Lambda\}$ and the Liouville form of $\left\{\mathrm{R},\,M;\,P_{\mathrm{R}},\,P_M\right\}$ is an exact form, which implies that the transformation of variables given by the set of equations (175) is canonical.

Armed with the certainty of the canonicity of the new variables, we can write the asymptotic form of the canonical variables and of the metric function $F(t,r)$. These are, for $r\to 0$

$$\begin{aligned}
F(t,r) &= 4R_2(t)\Lambda_0(t)^{-2}r^2+O(r^4)\,, && (178)\\
\mathrm{R}(t,r) &= R_0(t)+R_2(t)\,r^2+O(r^4)\,, && (179)\\
M(t,r) &= \frac{1}{32}\alpha^2R_0(t)^3+\frac{1}{32}\alpha R_0(t)R_2(t)\left(3\alpha^2R_0(t)-64R_2(t)\Lambda_0(t)^{-2}\right)\,r^2+O(r^4)\,, && (180)\\
P_{\mathrm{R}}(t,r) &= O(r)\,, && (181)\\
P_M(t,r) &= O(r)\,. && (182)
\end{aligned}$$

For $r\to\infty$, we have

$$\begin{aligned}
F(t,r) &= \frac{\alpha^2}{16}\,r^2-2(\eta(t)+2\rho(t))\,r^{-1}+O^\infty(r^{-2})\,, && (183)\\
\mathrm{R}(t,r) &= r+16\rho(t)\alpha^{-2}r^{-2}+O^\infty(r^{-3})\,, && (184)\\
M(t,r) &= M_+(t)+O^\infty(r^{-1})\,, && (185)\\
P_{\mathrm{R}}(t,r) &= O^\infty(r^{-4})\,, && (186)\\
P_M(t,r) &= O^\infty(r^{-6})\,, && (187)
\end{aligned}$$

where $M_+(t)=\alpha(\eta(t)+3\rho(t))$, as seen before in the surface terms (see Eq. (157)).

We are now almost ready to write the action with the new canonical variables. It is now necessary to determine the new Lagrange multipliers. In order to write the new constraints with the new Lagrange multipliers, we can use the identity given by the space derivative of M,

$$M'=-\Lambda^{-1}\left(R'H+\alpha^{-1}R^{-1}P_\Lambda H_r\right)\,. \qquad (188)$$

Solving for H and making use of the inverse transformations of Λ and P_Λ, in Eqs. (173) and (174), we get

$$\begin{aligned}
H &= -\frac{M'F^{-1}\mathrm{R}'+FP_MP_{\mathrm{R}}}{\left(F^{-1}(\mathrm{R}')^2-FP_M^2\right)^{\frac{1}{2}}}\,, && (189)\\
H_r &= P_MM'+P_{\mathrm{R}}\mathrm{R}'\,. && (190)
\end{aligned}$$

Following Kuchař [11], the new set of constraints, totally equivalent to the old set $H(t,r)=0$ and $H_r(t,r)=0$ outside the horizon, is $M'(t,r)=0$ and $P_{\mathrm{R}}(t,r)=0$. By continuity, this also applies on the horizon, where $F(t,r)=0$. So we can say that the equivalence is valid everywhere. So, the new Hamiltonian, the total sum of the constraints, can now be written as

$$NH + N^r H_R = N^M M' + N^{\mathrm{R}} P_{\mathrm{R}}\,. \tag{191}$$

In order to determine the new Lagrange multipliers, one has to write the left hand side of the previous equation, Eq. (191), and replace the constraints on that side by their expressions as functions of the new canonical coordinates, spelt out in Eqs. (189)-(190). After manipulation, one gets

$$\begin{aligned} N^M &= -\frac{NF^{-1}R'}{\left(F^{-1}(\mathrm{R}')^2 - FP_M^2\right)^{\frac{1}{2}}} + N^r P_M\,, && (192)\\ N^{\mathrm{R}} &= -\frac{NFP_M}{\left(F^{-1}(\mathrm{R}')^2 - FP_M^2\right)^{\frac{1}{2}}} + N^r R'\,. && (193)\end{aligned}$$

Using the inverse transformations Eqs. (173)-(174), and the identity $R=\mathrm{R}$, we can write the new multipliers as functions of the old variables

$$\begin{aligned} N^M &= -NF^{-1}R'\Lambda^{-1} + \alpha^{-1}N^rF^{-1}R^{-1}\Lambda P_\Lambda\,, && (194)\\ N^{\mathrm{R}} &= -\alpha^{-1}NR^{-1}P_\Lambda + N^r R'\,, && (195)\end{aligned}$$

allowing us to determine its asymptotic conditions from the original conditions given above. These transformations are non-singular for $r>0$. As before, for $r\to 0$,

$$\begin{aligned} N^M(t,r) &= -\frac{1}{2}N_1(t)\Lambda_0(t)R_2(t)^{-1} + O(r^2)\,, && (196)\\ N^{\mathrm{R}}(t,r) &= -2N_1^r(t)R_2(t)\,r^2 + O(r^4) = O(r^2)\,, && (197)\end{aligned}$$

and for $r\to\infty$ we have

$$\begin{aligned} N^M(t,r) &= -\tilde{N}_+(t) + O^\infty(r^{-4})\,, && (198)\\ N^{\mathrm{R}}(t,r) &= O^\infty(r^{-2})\,. && (199)\end{aligned}$$

These conditions (196)-(199) show that the transformations in Eqs. (194)-(195) are satisfactory in the case of $r\to\infty$, but not for $r\to 0$. This is due to fact that in order to fix the Lagrange multipliers for $r\to\infty$, as we are free to do, we fix $\tilde{N}_+(t)$, which we already do when adding the surface term

$$-\int dt\,\tilde{N}_+ M_+ \tag{200}$$

to the action, in order to obtain the equations of motion in the bulk, without surface terms. However, at $r=0$, we see that fixing the multiplier N^M to values independent of the canonical variables is not the same as fixing $N_1\Lambda_0^{-1}$ to values independent of the canonical variables. We need to rewrite the multiplier N^M for the asymptotic regime $r\to 0$ without affecting its behavior for $r\to\infty$. In order to proceed we have to make one assumption, which is that the expression given in asymptotic condition of $M(t,r)$, as $r\to 0$, for the term of order zero, $M_0 \equiv \frac{1}{32}\alpha^3 R_0(t)^3$, defines R_0 as a function of M_0, and R_0 is the horizon radius function, $R_0 \equiv R_{\mathrm{h}}(M_0)$. Also, we assume that $M_0>0$. With these assumptions, we are working in the domain of the classical solutions. We can immediately obtain that the variation of R_0 is given in relation to the variation of M_0 as

$$\delta R_0 = \frac{32}{3}\alpha^{-3}R_0^{-2}\delta M_0\,. \tag{201}$$

This expression will be used when we derive the equations of motion from the new action. We now define the new multiplier $\tilde{N}^M$ as

$$\tilde{N}^M = -N^M\left[(1-g) + 2g(\alpha R_0)\left(\frac{3\alpha^3}{16}R_0^2\right)^{-1}\right]^{-1}, \tag{202}$$

where $g(r) = 1 + O(r^2)$ for $r \to 0$ and $g(r) = O^\infty(r^{-5})$ for $r \to \infty$. This new multiplier, function of the old multiplier N^M, has as its properties for $r \to \infty$

$$\tilde{N}^M(t,r) = \tilde{N}_+(t) + O^\infty(r^{-5})\,, \tag{203}$$

and as its properties for $r \to 0$

$$\tilde{N}^M(t,r) = \tilde{N}_0^M(t) + O(r^2)\,, \tag{204}$$

where $\tilde{N}_0^M$ is given by

$$\tilde{N}_0^M = \frac{3}{64}\alpha^2 N_1 R_0 R_2^{-1} \Lambda_0\,. \tag{205}$$

When the constraint $M' = 0$ holds, the last expression is

$$\tilde{N}_0^M = N_1 \Lambda_0^{-1}\,. \tag{206}$$

With this new constraint $\tilde{N}^M$, fixing $N_1\Lambda_0^{-1}$ at $r = 0$ or fixing $\tilde{N}^M$ is equivalent, there being no problems with N^{R}, which is left as determined in Eq. (193).

The new action is now written as the sum of S_Σ, the bulk action, and $S_{\partial\Sigma}$, the surface action,

$$\begin{aligned} S\left[M, \mathrm{R}, P_M, P_{\mathrm{R}}; \tilde{N}^M, N^{\mathrm{R}}\right] &= \int dt \int_0^\infty dr \left(P_M \dot{M} + P_{\mathrm{R}}\dot{\mathrm{R}} - N^{\mathrm{R}} P_{\mathrm{R}} + \tilde{N}^M \left[(1-g) + 2gR_0 \left(\frac{3\alpha^2}{16} R_0^2\right)^{-1}\right] M'\right) \\ &+ \int dt \left(\frac{1}{2}\alpha R_0^2 \tilde{N}_0^M - \tilde{N}_+ M_+\right)\,. \end{aligned} \tag{207}$$

The new equations of motion are now

$$\begin{aligned} \dot{M} &= 0\,, && (208)\\ \dot{\mathrm{R}} &= N^{\mathrm{R}}\,, && (209)\\ \dot{P}_M &= (N^M)'\,, && (210)\\ \dot{P}_{\mathrm{R}} &= 0\,, && (211)\\ M' &= 0\,, && (212)\\ P_{\mathrm{R}} &= 0\,. && (213) \end{aligned}$$

Here we understood N^M to be a function of the new constraint, defined through Eq. (202). The resulting boundary terms of the variation of this new action, Eq. (207), are, first, terms proportional to δM and $\delta\mathrm{R}$ on the initial and final hypersurfaces, and, second, the term $\int dt \left(\frac{1}{2}\alpha R_0^2 \delta\tilde{N}_0^M - M_+\delta\tilde{N}_+\right)$. Here we have used the expression in Eq. (201). The action in Eq. (207) yields the equations of motion, Eqs. (208)-(213), provided that we fix the initial and final values of the new canonical variables and that we also fix the values of $\tilde{N}_0^M$ and of $\tilde{N}_+$. Thanks to the redefinition of the Lagrange multiplier, from N^M to $\tilde{N}^M$, the fixation of those quantities, $\tilde{N}_0^M$ and $\tilde{N}_+$, has the same meaning it had before the canonical transformations and the redefinition of N^M. This same meaning is guaranteed through the use of our gauge freedom to choose the multipliers, and at the same time not fixing the boundary variations independently of the choice of Lagrange multipliers, which in turn allow us to have a well defined variational principle for the action.

D. Hamiltonian reduction

We now solve the constraints in order to reduce to the true dynamical degrees of freedom. The equations of motion (208)-(213) allow us to write M as an independent function of the radial coordinate r,

$$M(t,r) = \mathbf{m}(t)\,. \tag{214}$$

The reduced action, with the constraints and Eq. (214) taken into account, is

$$S\left[\mathbf{m}, \mathbf{p_m}; \tilde{N}_0^M, \tilde{N}_+\right] = \int dt\, \mathbf{p_m}\dot{\mathbf{m}} - \mathbf{h}\,, \tag{215}$$

where

$$\mathbf{p_m} = \int_0^\infty dr\, P_M\,, \tag{216}$$

and the reduced Hamiltonian, $\mathbf{h}$, is now written as

$$\mathbf{h}(\mathbf{m};t) = -\frac{1}{2}\alpha R_{\rm h}^2 \tilde{N}_0^M + \tilde{N}_+\mathbf{m}\,, \tag{217}$$

with $R_{\rm h}$ being the horizon radius. We also have that $\mathbf{m} > 0$, according to the assumptions made in the previous subsection. Thanks to the functions $\tilde{N}_0^M(t)$ and $\tilde{N}_+(t)$ the Hamiltonian $\mathbf{h}$ is an explicitly time dependent function. The variational principle associated with the reduced action, Eq. (215), will fix the value of $\mathbf{m}$ on the initial and final hypersurfaces, or in the spirit of the classical analytical mechanics, the Hamiltonian principle fixes the initial and final values of the canonical coordinate. The equations of motion are

$$\dot{\mathbf{m}} \;=\; 0\,, \tag{218}$$

$$\dot{\mathbf{p}}_{\mathbf{m}} \;=\; \frac{32}{3}\,\alpha^{-2}R_{\rm h}^{-1}\tilde{N}_0^M - \tilde{N}_+\,. \tag{219}$$

The equation of motion for $\mathbf{m}$, Eq. (218), is understood as saying that $\mathbf{m}$ is, on a classical solution, equal to the mass parameter M of the solutions in Eq. (131). In order to interpret the other equation of motion, Eq. (218), we have to recall that from Eq. (171) one has $P_M = -T'$, where T is the Killing time. This, together with the definition of $\mathbf{p_m}$, given in Eq. (216), yields

$$\mathbf{p_m} = T_0 - T_+\,, \tag{220}$$

where T_0 is the value of the Killing time at the left end of the hypersurface of a certain t, and T_+ is the Killing time at spatial infinity, the right end of the same hypersurface of t. As the hypersurface evolves in the spacetime of the black hole solutions, the right hand side of Eq. (219) is equal to $\dot{T}_0 - \dot{T}_+$.

E. Quantum theory and partition function

The next step is to quantize the reduced Hamiltonian theory, by building the time evolution operator quantum mechanically and then obtaining a partition function through the analytic continuation of the same operator [12]-[16]. The variable $\mathbf{m}$ is regarded here as a configuration variable. This variable satisfies the inequality $\mathbf{m} > 0$. The wave functions will be of the form $\psi(\mathbf{m})$, with the inner product

$$(\psi,\chi) = \int_A \mu d\mathbf{m}\,\bar{\psi}\chi\,, \tag{221}$$

where A is the domain of integration defined by $\mathbf{m} > 0$ and $\mu(\mathbf{m})$ is a smooth and positive weight factor for the integration measure. It is assumed that μ is a slow varying function, otherwise arbitrary. We are thus working in the Hilbert space defined as $\mathscr{H} := L^2(A;\mu d\mathbf{m})$.

Again, the Hamiltonian operator, written as $\hat{\mathbf{h}}(t)$, acts through pointwise multiplication by the function $\mathbf{h}(\mathbf{m};t)$, which on a function of our working Hilbert space reads

$$\hat{\mathbf{h}}(t)\psi(\mathbf{m}) = \mathbf{h}(\mathbf{m};t)\psi(\mathbf{m})\,. \tag{222}$$

This Hamiltonian operator is an unbounded essentially self-adjoint operator. The corresponding time evolution operator in the same Hilbert space, which is unitary due to the fact that the Hamiltonian operator is self-adjoint, is

$$\hat{K}(t_2;t_1) = \exp\left[-i\int_{t_1}^{t_2} dt'\,\hat{\mathbf{h}}(t')\right]\,. \tag{223}$$

This operator acts also by pointwise multiplication in the Hilbert space. We define again

$$\mathcal{T} \;:=\; \int_{t_1}^{t_2} dt\,\tilde{N}_+(t)\,, \tag{224}$$

$$\Theta \;:=\; \int_{t_1}^{t_2} dt\,\tilde{N}_0^M(t)\,. \tag{225}$$

Using (217), (223), (224), and (225), we write the function which is in fact the action of the operator in the Hilbert space

$$K\left(\mathbf{m};\mathcal{T},\Theta\right)=\exp\left[-i\mathbf{m}\mathcal{T}+\frac{i}{2}\alpha R_{\mathrm{h}}^{2}\Theta\right]\,. \tag{226}$$

This expression indicates that $\hat{K}(t_2;t_1)$ depends on t_1 and t_2 only through the functions $\mathcal{T}$ and Θ. We can now write the operator corresponding to the function K as $\hat{K}(\mathcal{T};\Theta)$. The composition law in time $\hat{K}(t_3;t_2)\hat{K}(t_2;t_1)=\hat{K}(t_3;t_1)$ can now be regarded as a sum of the parameters inside $\hat{K}(\mathcal{T};\Theta)$. These parameters are evolution parameters defined by the boundary conditions, i.e., $\mathcal{T}$ is the Killing time elapsed at right spatial infinity and Θ is the boost parameter elapsed at the bifurcation circle.

F. Thermodynamics

We can now build the partition function for this system. The path to follow is to continue the operator to imaginary time and take the trace over a complete orthogonal base. Our classical thermodynamic situation consists of a three-dimensional circularly symmetric black hole with a particular dilaton function, asymptotically anti-de Sitter, in thermal equilibrium with a bath of Hawking radiation. Ignoring back reaction from the radiation, the geometry is described by the solution in Eqs. (131)-(132). Thus, we consider a thermodynamic ensemble in which the temperature, or more appropriately here, the inverse temperature β is fixed. This characterizes a canonical ensemble, and the partition function $\mathcal{Z}(\beta)$ arises naturally in such an ensemble. To analytically continue the Lorentzian solution to imaginary time, we put $\mathcal{T}=-i\beta$ and $\Theta=-2\pi i$, based on the regularity of the classical Euclidean solution. We arrive then at the following expression for the partition function

$$\mathcal{Z}(\beta)=\mathrm{Tr}\left[\hat{K}(-i\beta,-2\pi i)\right]\,. \tag{227}$$

From Eq. (226) this is realized as

$$\mathcal{Z}(\beta)=\int_0^\infty \mu\,d\mathbf{m}\,\exp\left[-\beta\mathbf{m}+\pi\alpha R_{\mathrm{h}}^{2}\right]\langle\mathbf{m}|\mathbf{m}\rangle\,. \tag{228}$$

Since $\langle\mathbf{m}|\mathbf{m}\rangle$ is equal to $\delta(0)$, one has to regularize (228). Again, following the Louko-Whiting procedure [12]-[16], we have to regularize and normalize the operator $\hat{K}$ beforehand. This leads to

$$\mathcal{Z}_{\mathrm{ren}}(\beta)=\mathcal{N}\int_A \mu\,d\mathbf{m}\,\exp\left[-\beta\mathbf{m}+\pi\alpha R_{\mathrm{h}}^{2}\right]\,, \tag{229}$$

where $\mathcal{N}$ is a normalization factor and A is the domain of integration. Provided the weight factor μ is slowly varying compared to the exponential in Eq. (229), and using the fact that the horizon radius R_{h} is a function of $\mathbf{m}$, the integral in Eq. (229) is convergent. Thermodynamically, $\mathbf{m}$ is analogous to the energy of the system. Changing integration variables, from $\mathbf{m}$ to R_{h}, where

$$\mathbf{m}=2^{-5}\alpha^3R_{\mathrm{h}}^{3}\,, \tag{230}$$

the integral Eq. (229) becomes

$$\mathcal{Z}_{\mathrm{ren}}(\beta)=\mathcal{N}\int_{A'}\widetilde{\mu}\,dR_{\mathrm{h}}\,\exp(-I_*)\,, \tag{231}$$

where A' is the new domain of integration after changing variables, and the function $I_*(R_{\mathrm{h}})$, a kind of an effective action (see [4]), is written as

$$I_*(R_{\mathrm{h}}):=2^{-5}\beta\,\alpha^3R_{\mathrm{h}}^{3}-\pi\alpha R_{\mathrm{h}}^{2}\,. \tag{232}$$

The domain of integration, A', is defined by the inequality $R_{\mathrm{h}}\geq 0$. The new weight factor $\widetilde{\mu}$ includes the Jacobian of the change of variables, which amounts to $\partial\mathbf{m}/\partial R_{\mathrm{h}}$. Since the weight factor is slowly varying, we can estimate the integral of $\mathcal{Z}_{\mathrm{ren}}(\beta)$ by the saddle point approximation. For that we have to calculate the critical points, which in our case amounts to finding the values of $R_{\rm h}$ for which the first derivative of $I_*(R_{\rm h})$ with respect to $R_{\rm h}$ is null. It happens for two different values of the domain

$$R_{\rm h}^- = 0\,, \tag{233}$$

$$R_{\rm h}^+ = 2^6\pi\left(3\beta\,\alpha^2\right)^{-1}\,. \tag{234}$$

In order to find the local extrema we evaluate the second derivative at these two points. One finds

$$\left.\frac{\partial^2 I_*}{\partial R_{\rm h}^2}\right|_{R_{\rm h}^-} = -2\pi\alpha\,, \tag{235}$$

$$\left.\frac{\partial^2 I_*}{\partial R_{\rm h}^2}\right|_{R_{\rm h}^+} = 2\pi\alpha\,. \tag{236}$$

Our domain starts at $R_{\rm h} = 0$, which is a local maximum. The global extremum, which is a minimum, is located at $R_{\rm h} = R_{\rm h}^+$. Evaluating the action I_* at $R_{\rm h}^+$ one obtains

$$I_*(R_{\rm h}^+) = 2^{-5}\beta\,\alpha^3(R_{\rm h}^+)^3 - \pi\alpha(R_{\rm h}^+)^2\,. \tag{237}$$

Substituting Eq. (234) into Eq. (237) gives

$$I_*(R_{\rm h}^+) = -2^{12}3^{-3}\pi^3\beta^{-2}\alpha^{-3}\,. \tag{238}$$

From Eq. (238) one sees that $I_*(R_{\rm h}^+) < 0$. By Taylor expanding the action in the exponential of the integral, Eq. (231), i.e.,

$$I_*(R_{\rm h}) = I_*(R_{\rm h}^+) + \left.\frac{\partial I_*}{\partial R_{\rm h}}\right|_{R_{\rm h}^+} R_{\rm h} + \frac{1}{2}\left.\frac{\partial^2 I_*}{\partial R_{\rm h}^2}\right|_{R_{\rm h}^+}(R_{\rm h})^2 + O((R_{\rm h})^3)\,, \tag{239}$$

we can separate the terms in such a way that we obtain the following expression for the renormalized partition function

$$\mathcal{Z}_{\rm ren}(\beta) = \exp\left[-I_*(R_{\rm h}^+)\right]\mathcal{N}\int_{A'}\widetilde{\mu}\,dR_{\rm h}\exp\left[-\pi\alpha R_{\rm h}^2\right]\,. \tag{240}$$

The Taylor expansion is up to second order, and evaluated at the critical point $R_{\rm h}^+$, which makes the first order term of the expansion of $I_*(R_{\rm h})$ disappear. The term which can be put outside the integral is the zero order term, which is the value of $I_*(R_{\rm h})$ at the extremum $R_{\rm h}^+$. The term left inside the exponential, $-\pi\alpha R_{\rm h}^2$, is minus the second order term in the Taylor expansion, where all the higher orders have been ignored, as this is a good approximation, provided the weight factor is slowly varying. Finally, we may write the renormalized partition function as

$$\mathcal{Z}_{\rm ren}(\beta) = {\rm P}\exp[2^{12}3^{-3}\pi^3\beta^{-2}\alpha^{-3}]\,, \tag{241}$$

where P is given by

$${\rm P} = \mathcal{N}\int_{A'}\widetilde{\mu}\,dR_{\rm h}\exp\left[-\pi\alpha R_{\rm h}^2\right]\,. \tag{242}$$

This P is a slowly varying prefactor and this approximation is better as we move to higher values of $|I_*(R_{\rm h}^+)|$. In the domain of integration the dominating contribution comes from the vicinity of $R_{\rm h} = R_{\rm h}^+$. Leaving the explicit dependence of the partition function on the $R_{\rm h}^+$ we write the logarithm of $\mathcal{Z}_{\rm ren}(\beta)$ as

$$\ln(\mathcal{Z}_{\rm ren}(\beta)) = \ln{\rm P} + \pi\,\alpha(R_{\rm h}^+)^2 - 2^{-5}\beta\,\alpha^3(R_{\rm h}^+)^3\,. \tag{243}$$

By ignoring the prefactor's logarithm, which closer to $R_{\rm h}^+$ is less relevant, we are able to determine the value of $\mathbf{m}$ at the critical point, where we find that it corresponds to the value of the mass of the classical solution of the black hole given in Eq. (131). Thus, when the critical point dominates the partition function, we have that the mean energy $\langle E\rangle$ is given by

$$\langle E\rangle = -\frac{\partial}{\partial\beta}\ln\mathcal{Z}_{\rm ren}(\beta) \approx 2^{-5}\alpha^3(R_{\rm h}^{+})^3 = \mathbf{m}^+ \,, \tag{244}$$

where $\mathbf{m}^+$ is obtained from Eq. (230) evaluated at $R_{\rm h}^+$. By replacing the value of $R_{\rm h}^+$ in Eq. (234) into Eq. (230), we write the temperature $\mathbf{T}\equiv\beta^{-1}$

$$\mathbf{T} = \left(\frac{3^{\frac{3}{2}}\,|\lambda|^3\,\mathbf{m}^+}{2^{\frac{5}{2}}\pi^3}\right)^{\frac{1}{3}} \,, \tag{245}$$

where $\mathbf{m}^+$ is the function in (230) evaluated at $R_{\rm h}^+$. Inverting Eq. (245) we obtain the function $\mathbf{m}^+(\beta)$, where β^{-1} is the inverse temperature,

$$\mathbf{m}^+(\beta) = 2^{13}\alpha^{-3}\pi^3(3\beta)^{-3}\,. \tag{246}$$

We see that $\partial\mathbf{m}^+/\partial\beta<0$, which through the heat capacity $C=-\beta^2(\partial\langle E\rangle/\partial\beta)$ tells us that the system is thermodynamically stable. The entropy is given by

$$S = \left(1-\beta\frac{\partial}{\partial\beta}\right)(\ln\mathcal{Z}_{\rm ren}(\beta)) \approx \pi\alpha {R_{\rm h}^{+}}^2 = 4\pi a {R_{\rm h}^{+}}^2\,. \tag{247}$$

This is the entropy of the three-dimensional black hole, which is a solution of the three-dimensional dilaton-gravity theory obtained through dimensionally reduced cylindrically general relativity (see also [26]). Note that the entropy obtained in Eq. (247) is not proportional to the area (i.e, circumference in the three-dimensional case) as the entropy of the BTZ black hole discussed in Section III. This is certainly due to the presence of the coupling of the dilaton to the metric.

V. HAMILTONIAN THERMODYNAMICS OF A REPRESENTATIVE DILATONIC BLACK HOLE ($\omega=-3$)

A. The metric

For $\omega=-3$, the general metric in Eq. (2) and the ϕ field in Eq. (6) reduce to the following

$$\begin{aligned} ds^2 &= -\left[(a\,R)^2-2\sqrt{a\,R}\,M\right]dT^2 + \frac{dR^2}{(a\,R)^2-2\sqrt{a\,R}\,M} + R^2\,d\varphi^2\,, && (248)\\ e^{-2\phi} &= \frac{1}{\sqrt{a\,R}}\,, && (249)\end{aligned}$$

with $2M=b$ and $a=\frac{4\sqrt{3}}{3}|\lambda|$. In Fig. 3 the Carter-Penrose diagram of the black hole solution for the case $\omega=-3$ is shown, which is analogous to the case $\omega=0$, where again the singularity at $R=0$ is a curvature singularity. Since we are now familiar with the whole formalism, we will be briefer in this section omitting several of the details.

B. Canonical formalism

With $\omega=-3$ the solution implies that $e^{-2\phi}=(a\,R)^{-\frac{1}{2}}$. The value of ω also implies that $a=\frac{4\sqrt{3}}{3}|\lambda|$ (see Eq. (8)). The action then becomes, up to surface terms,

$$\begin{aligned} S[\Lambda,\,R,\,\dot\Lambda,\,\dot R;\,N,\,N^r] \;=\; \int dt\int_0^\infty dr\,\Big\{ & 4\,(a\,R)^{-\frac{1}{2}}N\lambda^2\Lambda R-(a\,R)^{-\frac{1}{2}}N^{-1}\dot\Lambda\dot R+\frac{1}{4}(a\,R)^{-\frac{1}{2}}N^{-1}\Lambda R^{-1}\dot R^2\\ &+(a\,R)^{-\frac{1}{2}}N^{-1}\dot R(N^r\Lambda)'+(a\,R)^{-\frac{1}{2}}N^{-1}N^r\dot\Lambda R'-(a\,R)^{-\frac{1}{2}}N^{-1}N^r(N^r)'\Lambda R'\\ &-(a\,R)^{-\frac{1}{2}}N^{-1}(N^r)^2\Lambda' R'-\frac{1}{2}(a\,R)^{-\frac{1}{2}}N^{-1}N^r\Lambda R^{-1}\dot R R'+\frac{1}{4}(a\,R)^{-\frac{1}{2}}N^{-1}(N^r)^2\Lambda R^{-1}(R')^2\\ &+\frac{1}{4}(a\,R)^{-\frac{1}{2}}N\Lambda^{-1}R^{-1}(R')^2-(a\,R)^{-\frac{1}{2}}N(\Lambda^{-1})'R'-(a\,R)^{-\frac{1}{2}}N\Lambda^{-1}R''\Big\}\,. \end{aligned} \tag{250}$$



FIG. 3: The Carter-Penrose Diagram for the $\omega = -3$ case.

Depending on the situation we use three different letters containing the same information but with slightly different numerical values. Thus, a, λ, and l are related by $a^2 = \frac{16}{3}\lambda^2 = \frac{8}{3}l^{-2}$, where l is defined now as the AdS length. From the action above, we obtain the conjugate momenta

$$P_\Lambda = -N^{-1}(a\,R)^{-\frac{1}{2}}\left(\dot{R} - R'N^r\right), \tag{251}$$

$$P_R = -N^{-1}(a\,R)^{-\frac{1}{2}}\left\{\dot{\Lambda} - (\Lambda N^r)' - \frac{1}{2}\Lambda R^{-1}(\dot{R} - R'N^r)\right\}. \tag{252}$$

By performing a Legendre transformation we obtain the Hamiltonian, which is a sum of constraints, i.e.,

$$\begin{aligned}\mathcal{H} = {} & N\left\{-(a\,R)^{\frac{1}{2}}P_\Lambda P_R + \frac{1}{4}a^{\frac{1}{2}}R^{-\frac{1}{2}}\Lambda P_\Lambda^2 + (a\,R)^{-\frac{1}{2}}\left[(\Lambda^{-1})'R' + \Lambda^{-1}R'' - \frac{1}{4}\Lambda^{-1}(R')^2R^{-1}\right] - 4(a\,R)^{-\frac{1}{2}}\lambda^2\Lambda R\right\} \\ & + N^r\left\{P_R R' + \Lambda(P_\Lambda)'\right\} \equiv NH + N^r H_r\,. \end{aligned} \tag{253}$$

We can now write the action in Hamiltonian form, which reads

$$S[\Lambda,\, R,\, P_\Lambda,\, P_R;\, N,\, N^r] = \int dt \int_0^\infty dr\left\{P_\Lambda\dot{\Lambda} + P_R\dot{R} - NH - N^rH_r\right\}, \tag{254}$$

with the constraints defined in Eq. (253). From here we derive the equations of motion for the canonical variables and respective canonical momenta

$$\dot{\Lambda} = -N(a\,R)^{\frac{1}{2}}P_R + \frac{1}{2}Na^{\frac{1}{2}}R^{-\frac{1}{2}}\Lambda P_\Lambda + (N^r\Lambda)'\,, \tag{255}$$

$$\dot{R} = -N(a\,R)^{\frac{1}{2}}P_\Lambda + N^rR'\,, \tag{256}$$

$$\dot{P}_\Lambda = -\frac{1}{4}Na^{\frac{1}{2}}R^{-\frac{1}{2}}P_\Lambda^2 - \left(N(a\,R)^{-\frac{1}{2}}\right)'R'\Lambda^{-2} - \frac{1}{4}N(a\,R)^{-\frac{1}{2}}(R')^2R^{-1}\Lambda^{-2} + 4N(a\,R)^{-\frac{1}{2}}\lambda^2R + N^rP_\Lambda'\,, \tag{257}$$

$$\begin{aligned}\dot{P}_R = {} & \frac{1}{2}Na^{\frac{1}{2}}R^{-\frac{1}{2}}P_\Lambda P_R + \frac{1}{8}Na^{\frac{1}{2}}R^{-\frac{3}{2}}\Lambda P_\Lambda^2 - Na^{-\frac{1}{2}}(\Lambda^{-1})'(R^{-\frac{1}{2}})' \\ & + \frac{1}{4}N(a\,R)^{-\frac{1}{2}}\Lambda^{-1}R^{-1}R'' - \left(\left(Na^{-\frac{1}{2}}\right)'\Lambda^{-1}R^{-\frac{1}{2}}\right)' - \frac{1}{2}Na^{-\frac{1}{2}}\Lambda^{-1}(R^{-\frac{1}{2}})'' \\ & + 2Na^{-\frac{1}{2}}\lambda^2\Lambda R^{-\frac{1}{2}} + (N^rP_R)'\,. \end{aligned} \tag{258}$$

For a correct variational principle to be applied, we have to find out what surface terms are left over from the variation performed, with the purpose of deriving the equations of motion in Eqs. (255)-(258). These surface terms are,

$$\begin{aligned}\text{Surface terms} = {} & N(a\,R)^{-\frac{1}{2}}R'\Lambda^{-2}\delta\Lambda - N(a\,R)^{-\frac{1}{2}}\Lambda^{-1}\delta R' + \left((a\,R)^{-\frac{1}{2}}\right)'\Lambda^{-1}\delta R + \frac{1}{2}(a\,R)^{-\frac{1}{2}}\Lambda^{-1}R^{-1}R'\delta R \\ & - N^rP_R\delta R + N^r\Lambda\delta P_\Lambda\big|_0^\infty\,. \end{aligned} \tag{259}$$

In order to know the form of (259) for $r \to 0$, we assume

$$\begin{aligned}
\Lambda(t,r) &= \Lambda_0 + O(r^2)\,, && (260)\\
R(t,r) &= R_0 + R_2 r^2 + O(r^4)\,, && (261)\\
P_\Lambda(t,r) &= O(r^3) && (262)\\
P_R(t,r) &= O(r)\,, && (263)\\
N(t,r) &= N_1(t) r + O(r^3) && (264)\\
N^r(t,r) &= O(r^3)\,. && (265)
\end{aligned}$$

The surface terms become

$$\text{Surface terms}\big|_{r=0} = -N_1 (a\,R)^{-\frac{1}{2}} \Lambda_0^{-1} \delta R_0\,. \tag{266}$$

The asymptotic conditions for $r \to \infty$ are assumed as

$$\begin{aligned}
\Lambda(t,r) &= \frac{\sqrt{6}}{4}\, l r^{-1} + 2^{\frac{5}{4}} l^{\frac{5}{2}} \eta(t) r^{-\frac{5}{2}} + O^\infty(r^{-3})\,, && (267)\\
R(t,r) &= r + 2^{\frac{3}{4}} l^{\frac{3}{2}} \rho(t) r^{-\frac{1}{2}} + O^\infty(r^{-1})\,, && (268)\\
P_\Lambda(t,r) &= O^\infty(r^{-2}) && (269)\\
P_R(t,r) &= O^\infty(r^{-4})\,, && (270)\\
N(t,r) &= R(t,r)' \Lambda(t,r)^{-1} (\tilde{N}_+(t) + O^\infty(r^{-5}))\,, && (271)\\
N^r(t,r) &= O^\infty(r^{-2})\,, && (272)
\end{aligned}$$

where l is the AdS length. The surface terms, Eq. (259), for $r \to \infty$, are written as

$$\text{Surface terms}\big|_{r\to\infty} = \tilde{N}_+ \delta M_+\,, \tag{273}$$

where

$$M_+(t) = 2^5 3^{-\frac{5}{4}}\, \eta(t) + 2^2 3^{\frac{1}{2}}\, \rho(t)\,. \tag{274}$$

Therefore, the surface term added to (254) is

$$S_{\partial\Sigma}\left[\Lambda, R; N\right] = \int dt \left(2a^{-1} (a\,R_0)^{\frac{1}{2}} N_1 \Lambda_0^{-1} - \tilde{N}_+ M_+ \right)\,. \tag{275}$$

With this surface term added we obtain a well defined variational principle. What remains after variation of the total action, Eq. (254) and Eq. (275), is

$$\int dt \left(2a^{-1} (a\,R_0)^{\frac{1}{2}} \delta(N_1 \Lambda_0^{-1}) - \delta\tilde{N}_+ M_+ \right)\,. \tag{276}$$

The surface terms coming from the variation of the total action disappear as the result of the fixation of $\delta(N_1 \Lambda_0^{-1})$ on the horizon, $r = 0$, and of $\tilde{N}_+(t)$ at infinity, $r \to \infty$.

C. Reconstruction, canonical transformation, and action

Repeating the steps of the two previous sections, we now give the main results concerning $\omega = -3$. The metric function is given by the expression

$$F = (a\,R)^2 - 2\sqrt{a\,R}\,M\,. \tag{277}$$

The Killing time $T = T(t,r)$ is a function of (t,r), where we find that

$$-\,T' = F^{-1} \Lambda P_\Lambda (a\,R)^{\frac{1}{2}}\,, \tag{278}$$

which is equal to minus the conjugate momentum of the new variable M, or $P_M = P_M(t,r) \equiv -T'(t,r)$. With the help of Eq. (278) we find $F = F(t,r)$ as a function of the canonical variables,

$$F = (R')^2\Lambda^{-2} - aRP_\Lambda^2 \, . \tag{279}$$

Summing up, the canonical transformations are

$$\begin{aligned}
R &= \mathrm{R}\,, \\
M &= \frac{1}{2}(a\,R)^{-\frac{1}{2}}\left((a\,R)^2 - F\right)\,, \\
P_{\mathrm{R}} &= P_R - F^{-1}R^{-1}\Lambda^{-2}\left[RR'\Lambda P_\Lambda' + \frac{1}{2}(R')^2 P_\Lambda \Lambda - RR''\Lambda P_\Lambda + RR'\Lambda' P_\Lambda\right] \\
&\quad + \frac{1}{4}R^{-1}\Lambda P_\Lambda - 4\lambda^2 F^{-1} P_\Lambda R\Lambda\,, \\
P_M &= F^{-1}\Lambda P_\Lambda (a\,R)^{\frac{1}{2}}\,.
\end{aligned} \tag{280}$$

In addition, the relevant inverse transformations back to the old variable Λ and respective conjugate momentum P_Λ are

$$\Lambda = \left(F^{-1}(R')^2 - FP_M^2\right)^{\frac{1}{2}}\,, \tag{281}$$

$$P_\Lambda = (a\,R)^{-\frac{1}{2}} F P_M \left(F^{-1}(R')^2 - FP_M^2\right)^{-\frac{1}{2}}\,. \tag{282}$$

We have to show that this transformation is canonical. This requires using the identity

$$\begin{aligned}
P_\Lambda\delta\Lambda + P_R\delta R - P_M\delta M - P_{\mathrm{R}}\delta\mathrm{R} &= \left(\frac{1}{2}(a\,R)^{-\frac{1}{2}}\delta R \ln\left|\frac{(a\,R)^{-\frac{1}{2}}R' + \Lambda P_\Lambda}{(a\,R)^{-\frac{1}{2}}R' - \Lambda P_\Lambda}\right|\right)' + \\
&\quad + \delta\left(\Lambda P_\Lambda + \frac{1}{2}(a\,R)^{-\frac{1}{2}}R'\ln\left|\frac{(a\,R)^{-\frac{1}{2}}R' - \Lambda P_\Lambda}{(a\,R)^{-\frac{1}{2}}R' + \Lambda P_\Lambda}\right|\right)\,.
\end{aligned} \tag{283}$$

We now integrate Eq. (283), in r, from $r = 0$ to $r = \infty$. The first term on the right hand side of Eq. (283) vanishes due to the falloff conditions, Eqs. (260)-(265) and Eqs. (267)-(272). We obtain then the following expression

$$\int_0^\infty dr\ \left(P_\Lambda\delta\Lambda + P_R\delta R\right) - \int_0^\infty dr\ \left(P_M\delta M + P_{\mathrm{R}}\delta\mathrm{R}\right) = \delta\omega\left[\Lambda,\, R,\, P_\Lambda\right]\,, \tag{284}$$

where $\delta\omega\left[\Lambda,\, R,\, P_\Lambda\right]$ is a well defined functional, which is also an exact form. As above, this equality shows that the difference between the Liouville form of $\{R,\, \Lambda;\, P_R,\, P_\Lambda\}$ and the Liouville form of $\left\{\mathrm{R},\, M;\, P_{\mathrm{R}},\, P_M\right\}$ is an exact form, which implies that the set of transformations (280) is canonical.

With this result, we write the asymptotic conditions of the new canonical coordinates for $r \to 0$

$$F(t,r) = 4R_2^2\Lambda_0^{-2}r^2 + O(r^4)\,, \tag{285}$$

$$\mathrm{R}(t,r) = R_0 + R_2\,r^2 + O(r^4)\,, \tag{286}$$

$$M(t,r) = M_0 + M_2\,r^2 + O(r^4)\,, \tag{287}$$

$$P_{\mathrm{R}}(t,r) = O(r)\,, \tag{288}$$

$$P_M(t,r) = O(r)\,, \tag{289}$$

where we have

$$M_0 = 2\,3^{-\frac{3}{4}}l^{-2}R_0^2\left(\frac{\sqrt{2}}{2}l^{-1}R_0\right)^{-\frac{1}{2}}\,, \tag{290}$$

$$M_2 = -l^{-2}R_0R_2\left(\frac{\sqrt{2}}{2}l^{-1}R_0\right)^{\frac{1}{2}} + \frac{1}{4}\left(\frac{\sqrt{2}}{2}l^{-1}R_0\right)^{-\frac{1}{2}} 3^{\frac{1}{4}}\left[\frac{8}{3}l^{-2}R_0R_2 - 4R_2^2\Lambda_0^{-2}\right]\,. \tag{291}$$

For $r \to \infty$, we have

$$\begin{aligned} F(t,r) &= \frac{4}{3} l^{-2}\, r^2 - \frac{16}{3}\left(\frac{\sqrt{2}\, l^{-1}}{6}\right)^{\frac{1}{2}} \left(8\eta(t) + 3^{\frac{1}{2}}\rho(t)\right) r^{\frac{1}{2}} + O^\infty(r^0)\,, && (292) \\ \mathrm{R}(t,r) &= r + \left(\frac{\sqrt{2}}{2}\, l^{-1}\right)^{-\frac{3}{2}} \rho(t)\, r^{-\frac{1}{2}} + O^\infty(r^{-1})\,, && (293) \\ M(t,r) &= M_+(t) + O^\infty(r^{-\frac{1}{2}})\,, && (294) \\ P_{\mathrm{R}}(t,r) &= O^\infty(r^{-4})\,, && (295) \\ P_M(t,r) &= O^\infty(r^{-\frac{9}{5}})\,, && (296) \end{aligned}$$

where $M_+(t)$ is defined in Eq. (274).

We now write the future constraint M' as a function of the older constraints

$$M' = -\Lambda^{-1}\left(R'H + (a\,R)^{\frac{1}{2}} P_\Lambda H_r\right)\,. \qquad (297)$$

Using the inverse transformations of Λ and P_Λ in Eqs. (281) and (282), we obtain the same form for the old constraints as functions of the new variables

$$\begin{aligned} H &= -\frac{M' F^{-1} \mathrm{R}' + F P_M P_{\mathrm{R}}}{\left(F^{-1}(\mathrm{R}')^2 - F P_M^2\right)^{\frac{1}{2}}}\,, && (298) \\ H_r &= P_M M' + P_{\mathrm{R}} \mathrm{R}'\,. && (299) \end{aligned}$$

The new Hamiltonian, the total sum of the constraints, can now be written as

$$NH + N^r H_R = N^M M' + N^{\mathrm{R}} P_{\mathrm{R}}\,. \qquad (300)$$

The new multipliers are, using Eqs. (298)-(300),

$$\begin{aligned} N^M &= -\frac{N F^{-1} R'}{\left(F^{-1}(\mathrm{R}')^2 - F P_M^2\right)^{\frac{1}{2}}} + N^r P_M\,, && (301) \\ N^{\mathrm{R}} &= -\frac{N F P_M}{\left(F^{-1}(\mathrm{R}')^2 - F P_M^2\right)^{\frac{1}{2}}} + N^r R'\,. && (302) \end{aligned}$$

Using the inverse transformations Eqs. (281)-(282), and the identity $R = \mathrm{R}$, we can write the new multipliers as functions of the old variables

$$\begin{aligned} N^M &= -N F^{-1} R' \Lambda^{-1} + N^r F^{-1} \Lambda P_\Lambda (a\,R)^{\frac{1}{2}}\,, && (303) \\ N^{\mathrm{R}} &= -N P_\Lambda (a\,R)^{\frac{1}{2}} + N^r R'\,. && (304) \end{aligned}$$

For $r \to 0$ we have,

$$\begin{aligned} N^M(t,r) &= -\frac{1}{2} N_1(t) \Lambda_0 R_2^{-1} + O(r^2)\,, && (305) \\ N^{\mathrm{R}}(t,r) &= O(r^4)\,, && (306) \end{aligned}$$

and for $r \to \infty$ we have

$$\begin{aligned} N^M(t,r) &= -\tilde{N}_+(t) + O^\infty(r^{-2})\,, && (307) \\ N^{\mathrm{R}}(t,r) &= O^\infty(r^{-\frac{1}{2}})\,. && (308) \end{aligned}$$

Again, for $r \to 0$, fixing $N^M(t,r)$, which means fixing $N_1(t)\Lambda_0 R_2^{-1}$, is not equivalent to fixing $N_1 \Lambda_0^{-1}$. It is thus necessary to rewrite N^M for $r \to 0$. So, assuming M_0 as a function of R_0 allows one to define the horizon radius

$R_0 \equiv R_{\rm h}(M_0)$. We are thus working in the domain where $M_0 > 0$, the domain of the classical black hole solution. The variation of R_0 is given in terms of the variation of M_0 in the expression

$$\delta R_0 = \frac{4}{3} a^{-1} (a\, R)^{-\frac{1}{2}} \delta M_0 \,. \tag{309}$$

Remember that $a = \frac{4\sqrt{3}}{3}|\lambda|$. The new multiplier $\tilde{N}^M$ is obtained from the old N^M as

$$\tilde{N}^M = -N^M \left[(1-g) + \frac{1}{2} g l^2 R_0^{-1}\right]^{-1} , \tag{310}$$

where $l^2 = 2\lambda^2$, and $g(r) = 1 + O(r^2)$ for $r \to 0$ and $g(r) = O^\infty(r^{-5})$ for $r \to \infty$. This new multiplier, function of the old multiplier, $\tilde{N}^M$, has as its properties for $r \to \infty$

$$\tilde{N}^M(t,r) = \tilde{N}_+(t) + O^\infty(r^{-2}) \,, \tag{311}$$

and as its properties for $r \to 0$

$$\tilde{N}^M(t,r) = \tilde{N}_0^M(t) + O(r^2) \,, \tag{312}$$

where $\tilde{N}_0^M$ is given by

$$\tilde{N}_0^M = 2\lambda^2 N_1 R_0 R_2^{-1} \Lambda_0 \,. \tag{313}$$

When the constraint $M' = 0$ holds, the last expression is

$$\tilde{N}_0^M = N_1 \Lambda_0^{-1} \,. \tag{314}$$

With this new constraint $\tilde{N}^M$, fixing $N_1 \Lambda_0^{-1}$ or fixing $\tilde{N}^M$ is equivalent, there being no problem with $N^{\rm R}$, which is left as determined in Eq. (308).

The new action is then, summing both the bulk and the surface terms,

$$\begin{aligned} S\left[M, {\rm R}, P_M, P_{\rm R}; \tilde{N}^M, N^{\rm R}\right] \;=\; & \int dt \int_0^\infty dr \left(P_M \dot{M} + P_{\rm R} \dot{\rm R} - N^{\rm R} P_{\rm R} + \tilde{N}^M \left[(1-g) + \frac{1}{2} g\, l^2 \, R_0^{-1}\right] M'\right) + \\ & \int dt \left(2a^{-1}(a\, R_0)^{\frac{1}{2}} \tilde{N}_0^M - \tilde{N}_+ M_+\right) . \end{aligned} \tag{315}$$

The new equations of motion are now

$$\dot{M} \;=\; 0 \,, \tag{316}$$

$$\dot{\rm R} \;=\; N^{\rm R} \,, \tag{317}$$

$$\dot{P}_M \;=\; (N^M)' \,, \tag{318}$$

$$\dot{P}_{\rm R} \;=\; 0 \,, \tag{319}$$

$$M' \;=\; 0 \,, \tag{320}$$

$$P_{\rm R} \;=\; 0 \,. \tag{321}$$

Here we understood N^M to be a function of the new constraint, defined through Eq. (310). The resulting boundary terms of the variation of this new action, Eq. (315), are, first, terms proportional to δM and $\delta{\rm R}$ on the initial and final hypersurfaces, and, second, the term $\int dt \left(2a^{-1}(a\, R_0)^{\frac{1}{2}} \delta \tilde{N}_0^M - M_+ \delta \tilde{N}_+\right)$. Here we used the expression in Eq. (309). The action in Eq. (315) yields the equations of motion, Eqs. (316)-(321), provided that we fix the initial and final values of the new canonical variables and that we also fix the values of $\tilde{N}_0^M$ and of $\tilde{N}_+$. Thanks to the redefinition of the Lagrange multiplier, from N^M to $\tilde{N}^M$, the fixation of those quantities, $\tilde{N}_0^M$ and $\tilde{N}_+$, has the same meaning it had before the canonical transformations and the redefinition of N^M. This same meaning is guaranteed through the use of our gauge freedom to choose the multipliers, and at the same time not fixing the boundary variations independently of the choice of Lagrange multipliers, which in turn allow us to have a well defined variational principle for the action.

D. Hamiltonian reduction

We now solve the constraints in order to reduce to the true dynamical degrees of freedom. The equations of motion, Eqs. (316)-(321), allow us to write M as an independent function of the radial coordinate r,

$$M(t,r) = \mathbf{m}(t)\,. \tag{322}$$

The reduced action, with the constraints and Eq (322) taken into account, is

$$S\left[\mathbf{m}, \mathbf{p_m}; \tilde{N}_0^M, \tilde{N}_+\right] = \int dt\, \mathbf{p_m}\dot{\mathbf{m}} - \mathbf{h}\,, \tag{323}$$

where

$$\mathbf{p_m} = \int_0^\infty dr\, P_M\,, \tag{324}$$

and the reduced Hamiltonian, $\mathbf{h}$, is now written as

$$\mathbf{h}(\mathbf{m};t) = -2a^{-1}(a\,R_{\text{h}})^{\frac{1}{2}}\tilde{N}_0^M + \tilde{N}_+\mathbf{m}\,, \tag{325}$$

with R_{h} being the horizon radius. We also have that $\mathbf{m} > 0$. The equations of motion are then

$$\begin{aligned} \dot{\mathbf{m}} &= 0\,, && (326)\\ \dot{\mathbf{p}}_{\mathbf{m}} &= \frac{4}{3}a^{-2}(R_{\text{h}})^{-1}\tilde{N}_0^M - \tilde{N}_+\,. && (327) \end{aligned}$$

Here $\mathbf{m}$ is equal to the mass parameter M of the classical solution in Eq. (248). The second equation of motion, Eq. (327), describes the time evolution of the difference of the Killing times on the horizon and at infinity, due to $\mathbf{p_m} = T_0 - T_+$ and Eq. (324).

E. Quantum theory and partition function

The steps developed in Sections III and IV can be readily used here. So we do nont spell out the corresponding calculations in detail.

F. Thermodynamics

We can now build the partition function for this system, with the Hamiltonian given in Eq. (325). The steps are the same as was the case with $\omega = \infty$ and $\omega = 0$. The thermodynamic ensemble is also the canonical ensemble. Thus, the operator is

$$K\left(\mathbf{m}; \mathcal{T}, \Theta\right) = \exp\left[-i\mathbf{m}\mathcal{T} + 2\,i\,a^{-1}\,(a\,R_{\text{h}})^{\frac{1}{2}}\Theta\right]\,. \tag{328}$$

Again, with $\mathcal{T} = -i\beta$ and $\Theta = -2\pi i$, we write the general form of the partition function as

$$\mathcal{Z} = Tr\left[\hat{K}(-i\beta, -2\pi i)\right]\,. \tag{329}$$

This is realized as

$$\mathcal{Z}_{\text{ren}}(\beta) = \mathcal{N}\int_{A'} \widetilde{\mu}\, dR_{\text{h}}\, \exp(-I_*)\,, \tag{330}$$

where $\mathcal{N}$ is a normalization factor, A' is the domain of integration, and the function $I_*(R_{\text{h}})$, a kind of an effective action [4]. $I_*(R_{\text{h}})$, is written as

$$I_*(R_{\text{h}}) := \frac{\beta}{2}(a\,R_{\text{h}})^{\frac{3}{2}} - 4\pi a^{-1}(a\,R_{\text{h}})^{\frac{1}{2}}\,. \tag{331}$$

The domain of integration, given by A', is defined by the inequality $R_{\rm h} \geq 0$. The new weight factor $\widetilde{\mu}$ includes the Jacobian of the change of variables, which amounts to $\partial\mathbf{m}/\partial R_{\rm h}$. Its critical point is at $R_{\rm h}^{+} = \frac{8}{3}a^{-2}\pi\beta^{-1}$. The action evaluated at the critical point is

$$I_*(R_{\rm h}^{+}) = -a^{-\frac{3}{2}}\beta^{-\frac{1}{2}}\left(\frac{8}{3}\pi\right)^{\frac{3}{2}} . \tag{332}$$

From Eq. (332) one sees that $I_*(R_{\rm h}^{+}) < 0$. It is seen that the critical point is a minimum. This implies that

$$\begin{aligned} \mathcal{Z}_{\rm ren}(\beta) &\approx P\exp\left[-I_*(R_{\rm h}^{+})\right] \\ &= P\exp\left[-\frac{\beta}{2}(a\,R_{\rm h}^{+})^{\frac{3}{2}}\right] . \end{aligned} \tag{333}$$

We can now derive the basic thermodynamic results, as long as the approximation for the saddle point is valid, which means that we have to work in the neighborhood of the classical solution (248), where the critical point dominates. The expected value of the energy E is

$$\langle E\rangle = -\frac{\partial}{\partial\beta}\ln\mathcal{Z}_{\rm ren}(\beta) \approx \frac{1}{2}(a\,R_{\rm h}^{+})^{\frac{3}{2}} = \mathbf{m}^{+} . \tag{334}$$

Here we have

$$\mathbf{T} = \left(\frac{3^{\frac{3}{4}}\,|\lambda|^{\frac{3}{2}}\,\mathbf{m}^{+}}{2^{\frac{1}{2}}\pi^{\frac{3}{2}}}\right)^{\frac{2}{3}} , \tag{335}$$

$$\mathbf{m}^{+}(\beta) = \frac{1}{2}a^{-\frac{3}{2}}\beta^{-\frac{3}{2}}\left(\frac{8}{3}\pi\right)^{\frac{3}{2}} . \tag{336}$$

The derivative of $\mathbf{m}^{+}(\beta)$ with respect to β is negative, which means that the heat capacity is positive. The system is thus stable. Finally the entropy is given by

$$S = \left(1-\beta\frac{\partial}{\partial\beta}\right)(\ln\mathcal{Z}_{\rm ren}(\beta)) \approx 4\pi\sqrt{a^{-1}R_{\rm h}^{+}} . \tag{337}$$

The result recovers the entropy for the three-dimensional dilatonic black hole with $\omega = -3$ (see also [26]).

VI. CONCLUSIONS

We have continued the Louko-Whiting program of studying, through Hamiltonian methods, the thermodynamic properties of black holes in several theories in different dimensions (see [12] and [13]-[19]).

Specifically we have calculated the thermodynamic properties of black hole solutions with asymptotic infinities that allow a well formulated Hamiltonian formalism in three-dimensional Brans-Dicke dilaton-gravity. Only certain values of the Brans-Dicke parameter ω are allowed in this juncture. The corresponding theories are general relativity, i.e., $\omega \to \infty$, a dimensionally reduced cylindrical four-dimensional general relativity theory, i.e., $\omega = 0$, and a theory representing a class of dilaton-gravity theories, with a typical ω given by $\omega = -3$. Within a three-dimensional context, we have built a framework for the classical Hamiltonian theory, where the metric functions are used as canonical variables, and where one foliates the spacetime with equal time spacelike hypersurfaces. These hypersurfaces go from the bifurcation circle of the horizon on the left, to the asymptotic anti-de Sitter infinity on the right. Then we have performed a canonical transformation based on a reconstruction of the canonical variables. This canonical transformation is an adaptation of Kuchař's work [11] to the Louko-Whiting method [12, 13, 14, 15, 16]. In [11] the boundaries are the left and right infinities of the Kruskal diagram (or if one wishes of the Carter-Penrose diagram), whereas in [12, 13, 14, 15, 16] and here, with a thermodynamic goal in mind, the boundaries are the bifurcation sphere, here a 1-sphere or circle, and the right infinity. When the classical equations hold one finds that the new canonical coordinate is indeed the physical parameter of the classical solutions of the dilatonic black holes in three dimensions, i.e., the black hole mass M. Its conjugate momentum, $P_M = -T'$, is the spatial derivative of the Killing time. It survives the reduction, i.e., the elimination of constraints, becoming at last the difference between the Killing time at the bifurcation circle and at infinity. Its equation of motion is the equation for the evolution of the difference between Killing times, with respect to the time parameter t of each hypersurface. The other coordinate, R, and respective momentum, $P_{\rm R}$, vanish after reduction, being pure gauge. With the new variables, come new constraints and new Lagrange multipliers. We then reduce the Hamiltonian to an unconstrained theory with one pair of canonical coordinates. Then one performs a quantization of the theory, where one replaces the functions of the physical observables for operators of a Hilbert space. Constructing then the Schrödinger evolution operator from the Hamiltonian and taking the trace on a suitable basis, we obtain the partition function of the canonical thermodynamic ensemble. This ensemble is well defined and under suitable conditions the classical Euclidean solutions dominate the partition function, yielding the thermodynamics of the systems.

To sum up, as noted above, this formalism was previously applied to two [13] (see also [18, 19]), four [12, 14, 16, 17], and five dimensions [15] in several different theories. Here we have applied to three dimensions in a quite general dilaton-gravity theory. We have shown that in three-dimensional theories with well defined asymptotics the formalism fits well. As in other instances, the negative cosmological constant has had a stabilizing role to play here in what the thermodynamic results are concerned. Notwithstanding, several modifications were needed. First, in the powers of the fall-off conditions, and second due to the presence of the scalar dilaton field, which was reflected in the fact that it changed the powers of the radial R coordinate, to name a few. Although, in order to build a three-dimensional Lorentzian Hamiltonian theory, these modifications had to be made, we have derived a quantum theory and a statistical description of the systems in question, and found the corresponding thermodynamics, with precise values for the temperature and entropy of the black holes studied.

Acknowledgments

GASD is supported by grant SFRH/BD/2003 from FCT. This work was partially funded by Fundação para a Ciência e Tecnologia (FCT) - Portugal, through project POCI/FP/63943/2005.

[1] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
[2] J. D. Bekenstein, Phys. Rev. D **9**, 3292 (1974).
[3] S. W. Hawking, in *General Relativity: An Einstein Centenary Survey*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1979).
[4] J. W. York, Phys. Rev. D **33**, 2092 (1986).
[5] O. B. Zaslavskii, Phys. Lett. A **152**, 463 (1991).
[6] H. W. Braden, J. D. Brown, B. F. Whiting, and J. W. York, Phys. Rev. D **42**, 3376 (1990).
[7] C. S. Peça and J. P. S. Lemos, Phys. Rev. D **59**, 124007 (1999).
[8] P. A. M. Dirac, *Lectures on Quantum Mechanics*, (Yeshiva University, New York, 1964).
[9] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: an introduction to current research*, edited by L. Witten (Wiley, New York, 1962), Chap. 7 [gr-qc/0405109].
[10] T. Regge and C. Teitelboim, Ann. Phys. **88**, 286 (1974).
[11] K. V. Kuchař, Phys. Rev. D **50**, 3961 (1994).
[12] J. Louko and B. F. Whiting, Phys. Rev. D **51**, 5583 (1995).
[13] S. Bose, J. Louko, L. Parker, and Y. Peleg, Phys. Rev. D **53**, 5708 (1996).
[14] J. Louko and S. N. Winters-Hilt, Phys. Rev. D **54**, 2647 (1996).
[15] J. Louko, J. Z. Simon, and S. N. Winters-Hilt, Phys. Rev. D **55** 3525 (1997).
[16] D. R. Brill, J. Louko, and P. Peldan, Phys. Rev. D **56** 3600 (1997).
[17] S. Bose, L. Parker, and Y. Peleg, Phys. Rev. D **56**, 987 (1997).
[18] G. Kunstatter, R. Petryk, and S. Shelemy, Phys. Rev. D **57**, 3537 (1998).
[19] A. J. M. Medved and G. Kunstatter, Phys. Rev. D **59**, 104005 (1999).
[20] M. Bañados, C. Teitelboim, and J. Zanelli, Phys. Rev. Lett. **69**, 1849 (1992).
[21] M. Bañados, M. Henneaux, C. Teitelboim, and J. Zanelli, Phys. Rev. D **48**, 1506 (1993).
[22] G. T. Horowitz and D. L. Welch, Phys. Rev. Lett. **71** 328 (1993).
[23] J. Horne and G. T. Horowitz, Nucl. Phys. B **368**, 444 (1992).
[24] J. P. S. Lemos, Phys Lett. B **353**, 46 (1995).
[25] P. M. Sá, A. Kleber, and J. P. S. Lemos, Class. Quantum Grav. **13**, 125 (1996).
[26] M. Natsuume and T. Okamura, Phys. Rev. D **62**, 064027 (2000).
[27] O. B. Zaslavskii, Class. Quant. Grav. **11**, L33 (1994).
[28] S. Carlip, Class. Quantum Grav. **12**, 2853 (1995).